

TANNAKIAN DUALITY FOR AFFINE HOMOGENEOUS SPACES

TEODOR BANICA

ABSTRACT. Associated to any closed quantum subgroup $G \subset U_N^+$ and any index set $I \subset \{1, \dots, N\}$ is a certain homogeneous space $X_{G,I} \subset S_{\mathbb{C},+}^{N-1}$, called affine homogeneous space. We discuss here the abstract axiomatization of the algebraic manifolds $X \subset S_{\mathbb{C},+}^{N-1}$ which can appear in this way, by using Tannakian duality methods.

INTRODUCTION

The compact quantum groups were introduced by Woronowicz in [13], [14]. These are abstract objects, having no points in general, generalizing at the same time the usual compact groups, and the duals of the discrete groups. The compact quantum groups have no Lie algebra in general, but analogues of the Peter-Weyl theory, of Tannakian duality, and of the Weingarten integration formula, are available. Thus, we have here some interesting examples of noncommutative manifolds, which are definitely algebraic, and which are probably a bit Riemannian too, because we can integrate on them.

The present work is a continuation of [1], which was concerned with the integration theory over the associated homogeneous spaces. The main finding there was the fact that, in order to have a good integration theory, one must restrict the attention to a certain special class of homogeneous spaces, called “affine”. To be more precise, associated to any closed subgroup $G \subset U_N^+$ and any index set $I \subset \{1, \dots, N\}$ is a certain homogeneous space $X_{G,I} \subset S_{\mathbb{C},+}^{N-1}$, called affine. In the classical case this space appears as $X_{G,I} = G/(G \cap C_N^I)$, where $C_N^I \subset U_N$ is the group of unitaries fixing the vector $\xi_I = \frac{1}{\sqrt{|I|}} \sum_{i \in I} e_i$. In general, however, there are many new twists and questions coming from noncommutativity. Importantly, this construction covers many interesting examples. See [1].

One question left open in [1] was that of finding an abstract axiomatization of the algebraic manifolds $X \subset S_{\mathbb{C},+}^{N-1}$ which can appear in this way. We will answer here this question, with a Tannakian characterization of such manifolds. We believe that some further improvements of this result can lead to an axiomatization of the “easy algebraic manifolds”, which was the main question in [1], and which remains open.

The paper is organized as follows: 1 is a preliminary section, in 2-3 we state and prove the Tannakian duality result, and 4 contains a number of further results.

2010 *Mathematics Subject Classification.* 46L65 (46L89).

Key words and phrases. Quantum isometry, Noncommutative manifold.

1. HOMOGENEOUS SPACES

We use Woronowicz's quantum group formalism in [13], [14], with the extra axiom $S^2 = id$. To be more precise, the definition that we will need is:

Definition 1.1. *Assume that A is a unital C^* -algebra, and that $u \in M_N(A)$ is a unitary matrix whose coefficients generate A , such that the following formulae define morphisms of C^* -algebras $\Delta : A \rightarrow A \otimes A$, $\varepsilon : A \rightarrow \mathbb{C}$ and $S : A \rightarrow A^{opp}$:*

$$\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj} \quad , \quad \varepsilon(u_{ij}) = \delta_{ij} \quad , \quad S(u_{ij}) = u_{ji}^*$$

We write then $A = C(G)$, and call G a compact matrix quantum group.

The above maps Δ, ε, S are called comultiplication, counit and antipode. The basic examples include the compact Lie groups $G \subset U_N$, their q -deformations at $q = -1$, and the duals of the finitely generated discrete groups $\Gamma = \langle g_1, \dots, g_N \rangle$. See [13].

We recall that the free unitary quantum group U_N^+ , constructed by Wang in [11], and the corresponding free complex sphere $S_{\mathbb{C},+}^{N-1}$, from [3], are constructed as follows:

$$\begin{aligned} C(U_N^+) &= C^* \left((u_{ij})_{i,j=1,\dots,N} \mid u^* = u^{-1}, u^t = \bar{u}^{-1} \right) \\ C(S_{\mathbb{C},+}^{N-1}) &= C^* \left(x_1, \dots, x_N \mid \sum_i x_i x_i^* = \sum_i x_i^* x_i = 1 \right) \end{aligned}$$

Here both the algebras on the right are by definition universal C^* -algebras.

Following [1], we can now formulate the following definition:

Definition 1.2. *An affine homogeneous space over a closed subgroup $G \subset U_N^+$ is a closed subset $X \subset S_{\mathbb{C},+}^{N-1}$, such that there exists an index set $I \subset \{1, \dots, N\}$ such that*

$$\alpha(x_i) = \frac{1}{\sqrt{|I|}} \sum_{j \in I} u_{ij} \quad , \quad \Phi(x_i) = \sum_j u_{ij} \otimes x_j$$

define morphisms of C^* -algebras, satisfying $(\int_G \otimes id)\Phi = \int_G \alpha(\cdot)1$.

Here, and in what follows, a closed subspace $Y \subset Z$ corresponds by definition to a quotient map $C(Z) \rightarrow C(Y)$. As for \int_G , this is the Haar integration. See [13].

As a first observation, the coaction condition $(id \otimes \Phi)\Phi = (\Delta \otimes id)\Phi$ is satisfied, and we have as well $(id \otimes \alpha)\Phi = \Delta\alpha$. In the case where α is injective, we have:

Proposition 1.3. *When α is injective we must have $X = X_{G,I}^{min}$, where:*

$$C(X_{G,I}^{min}) = \left\langle \frac{1}{\sqrt{|I|}} \sum_{j \in I} u_{ij} \mid i = 1, \dots, N \right\rangle \subset C(G)$$

Moreover, $X_{G,I}^{min}$ is affine homogeneous, for any $G \subset U_N^+$, and any $I \subset \{1, \dots, N\}$.

Proof. The first assertion is clear from definitions. Regarding now the second assertion, consider the variables $z_i = \frac{1}{\sqrt{|I|}} \sum_{j \in I} u_{ij} \in C(G)$. We have then:

$$\Delta(z_i) = \frac{1}{\sqrt{|I|}} \sum_{j \in I} \sum_k u_{ik} \otimes u_{kj} = \sum_k u_{ik} \otimes z_k$$

Thus we have a coaction map as in Definition 1.2, given by $\Phi = \Delta$, and the ergodicity condition, namely $(\int_G \otimes id)\Delta = \int_G(\cdot)1$, holds as well, by definition of \int_G . See [1]. \square

Given exponents $e_1, \dots, e_k \in \{1, *\}$, consider the following quantities:

$$P_{i_1 \dots i_k, j_1 \dots j_k} = \int_G u_{i_1 j_1}^{e_1} \dots u_{i_k j_k}^{e_k}$$

Once again following [1], we have the following result:

Proposition 1.4. *We must have $X \subset X_{G,I}^{max}$, as subsets of $S_{\mathbb{C},+}^{N-1}$, where:*

$$C(X_{G,I}^{max}) = C(S_{\mathbb{C},+}^{N-1}) / \left\langle (Px^{\otimes k})_{i_1 \dots i_k} = \frac{1}{\sqrt{|I|^k}} \sum_{j_1 \dots j_k \in I} P_{i_1 \dots i_k, j_1 \dots j_k} \mid \forall k, \forall i_1, \dots, i_k \right\rangle$$

Moreover, $X_{G,I}^{max}$ is affine homogeneous, for any $G \subset U_N^+$, and any $I \subset \{1, \dots, N\}$.

Proof. The idea here is that the ergodicity condition $(\int_G \otimes id)\Phi = \int_G \alpha(\cdot)1$ produces the relations in the statement. To be more precise, observe that we have:

$$\begin{aligned} & \left(\int_G \otimes id \right) \Phi = \int \alpha(\cdot)1 \\ \iff & \left(\int_G \otimes id \right) \Phi(x_{i_1}^{e_1} \dots x_{i_k}^{e_k}) = \frac{1}{\sqrt{|I|^k}} \int_G \alpha(x_{i_1}^{e_1} \dots x_{i_k}^{e_k}), \forall k, \forall i_1, \dots, i_k \\ \iff & \sum_{j_1 \dots j_k} P_{i_1 \dots i_k, j_1 \dots j_k} x_{j_1}^{e_1} \dots x_{j_k}^{e_k} = \frac{1}{\sqrt{|I|^k}} \sum_{j_1 \dots j_k \in I} P_{i_1 \dots i_k, j_1 \dots j_k}, \forall k, \forall i_1, \dots, i_k \end{aligned}$$

Thus we have $X \subset X_{G,I}^{max}$, and the last assertion is standard as well. See [1]. \square

We will need one more general result from [1], namely an extension of the Weingarten integration formula [2], [5], [6], [12], to the affine homogeneous space setting:

Proposition 1.5. *Assuming that $G \rightarrow X$ is an affine homogeneous space, with index set $I \subset \{1, \dots, N\}$, the Haar integration functional $\int_X = \int_G \alpha$ is given by*

$$\int_X x_{i_1}^{e_1} \dots x_{i_k}^{e_k} = \sum_{\pi, \sigma \in D} (\xi_\pi)_{i_1 \dots i_k} K_I(\sigma) W_{kN}(\pi, \sigma)$$

where $\{\xi_\pi \mid \pi \in D\}$ is a basis of $Fix(u^{\otimes k})$, $W_{kN} = G_{kN}^{-1}$ with $G_{kN}(\pi, \sigma) = \langle \xi_\pi, \xi_\sigma \rangle$ is the associated Weingarten matrix, and $K_I(\sigma) = \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \overline{(\xi_\sigma)_{b_1 \dots b_k}}$.

Proof. By using the Weingarten formula for the quantum group G , we have:

$$\begin{aligned} \int_X x_{i_1}^{e_1} \dots x_{i_k}^{e_k} &= \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \int_G u_{i_1 b_1}^{e_1} \dots u_{i_k b_k}^{e_k} \\ &= \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \sum_{\pi, \sigma \in D} (\xi_\pi)_{i_1 \dots i_k} \overline{(\xi_\sigma)_{b_1 \dots b_k}} W_{kN}(\pi, \sigma) \end{aligned}$$

But this gives the formula in the statement, and we are done. See [1]. \square

Finally, here is a natural example of an intermediate space $X_{G,I}^{min} \subset X \subset X_{G,I}^{max}$:

Proposition 1.6. *Given a closed quantum subgroup $G \subset U_N^+$, and a set $I \subset \{1, \dots, N\}$, if we consider the following quotient algebra*

$$C(X_{G,I}^{med}) = C(S_{\mathbb{C},+}^{N-1}) / \left\langle \sum_{a_1 \dots a_k} \xi_{a_1 \dots a_k} x_{a_1}^{e_1} \dots x_{a_k}^{e_k} = \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \xi_{b_1 \dots b_k} \mid \forall k, \forall \xi \in \text{Fix}(u^{\otimes k}) \right\rangle$$

we obtain in this way an affine homogeneous space $G \rightarrow X_{G,I}$.

Proof. We know from Proposition 1.4 above that $X_{G,I}^{max} \subset S_{\mathbb{C},+}^{N-1}$ is constructed by imposing to the standard coordinates the conditions $Px^{\otimes k} = P^I$, where:

$$P_{i_1 \dots i_k, j_1 \dots j_k} = \int_G u_{i_1 j_1}^{e_1} \dots u_{i_k j_k}^{e_k} \quad , \quad P_{i_1 \dots i_k}^I = \frac{1}{\sqrt{|I|^k}} \sum_{j_1 \dots j_k \in I} P_{i_1 \dots i_k, j_1 \dots j_k}$$

According to the Weingarten integration formula for G , we have:

$$\begin{aligned} (Px^{\otimes k})_{i_1 \dots i_k} &= \sum_{a_1 \dots a_k} \sum_{\pi, \sigma \in D} (\xi_\pi)_{i_1 \dots i_k} \overline{(\xi_\sigma)_{a_1 \dots a_k}} W_{kN}(\pi, \sigma) x_{a_1}^{e_1} \dots x_{a_k}^{e_k} \\ P_{i_1 \dots i_k}^I &= \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \sum_{\pi, \sigma \in D} (\xi_\pi)_{i_1 \dots i_k} \overline{(\xi_\sigma)_{b_1 \dots b_k}} W_{kN}(\pi, \sigma) \end{aligned}$$

Thus $X_{G,I}^{med} \subset X_{G,I}^{max}$, and the other assertions are standard as well. See [1]. \square

We can now put everything together, as follows:

Theorem 1.7. *Given a closed subgroup $G \subset U_N^+$, and a subset $I \subset \{1, \dots, N\}$, the affine homogeneous spaces over G , with index set I , have the following properties:*

- (1) *These are exactly the intermediate subspaces $X_{G,I}^{min} \subset X \subset X_{G,I}^{max}$ on which G acts affinely, with the action being ergodic.*
- (2) *For the minimal and maximal spaces $X_{G,I}^{min}$ and $X_{G,I}^{max}$, as well as for the intermediate space $X_{G,I}^{med}$ constructed above, these conditions are satisfied.*
- (3) *By performing the GNS construction with respect to the Haar integration functional $\int_X = \int_G \alpha$ we obtain the minimal space $X_{G,I}^{min}$.*

We agree to identify all these spaces, via the GNS construction, and denote them $X_{G,I}$.

Proof. This follows indeed by combining the various results and observations formulated above. Once again, for full details on all these facts, we refer to [1]. \square

Observe the similarity with what happens for the C^* -algebras of the discrete groups, where the various intermediate algebras $C^*(\Gamma) \rightarrow A \rightarrow C_{red}^*(\Gamma)$ must be identified as well, in order to reach to a unique noncommutative space $\widehat{\Gamma}$. For details here, see [13].

Regarding now the basic examples of such spaces, we have:

Proposition 1.8. *Given $N \in \mathbb{N}$ and $I \subset \{1, \dots, N\}$, the following hold:*

- (1) *In the classical case, $G \subset U_N$, we have $X_{G,I} = G/(G \cap C_N^I)$, where $C_N^I \subset U_N$ is the group of unitaries fixing the vector $\xi_I = \frac{1}{\sqrt{|I|}}(\delta_{i \in I})_i$.*
- (2) *In the group dual case, $G = \widehat{\Gamma} \subset U_N^+$ with $\Gamma = \langle g_1, \dots, g_N \rangle$, embedded via $u_{ij} = \delta_{ij}g_i$, we have $X_{G,I} = \widehat{\Gamma}_I$, with $\Gamma_I = \langle g_i | i \in I \rangle \subset \Gamma$.*

Proof. In this statement (1) follows from the fact that the action $G \curvearrowright X_{G,I}$ can be shown to be transitive, and the stabilizer of ξ_I is the group $G \cap C_N^I$ in the statement. As for (2), this follows directly from Definition 1.2, by using $u_{ij} = \delta_{ij}g_i$. See [1]. \square

One interesting question is that of understanding how much of (1) can apply to the general case. The answer here is as follows, with (2) providing counterexamples:

Proposition 1.9. *We have a quotient map as follows, which is in general proper,*

$$G/(G \cap C_N^{I+}) \rightarrow X_{G,I}$$

where $C_N^{I+} \subset U_N^+$ is the subgroup defined by $C(C_N^{I+}) = C(U_N^+)/\langle u\xi_I = \xi_I \rangle$, with the relation $u\xi_I = \xi_I$ being interpreted as an equality of column vectors, over $C(U_N^+)$.

Proof. Observe first that C_N^{I+} is indeed a quantum group. In fact, it is standard to exhibit an isomorphism $C_N^{+I} \simeq U_{N-1}^+$, by reasoning as in [10]. We must check that the defining relations for $C(G/(G \cap C_N^{I+}))$ hold for the standard generators $x_i \in C(X_{G,I})$. But if we denote by $\pi : C(G) \rightarrow C(G \cap C_N^{I+})$ the quotient map, we have, as desired:

$$(id \otimes \pi)\Delta x_i = (id \otimes \pi) \left(\frac{1}{\sqrt{|I|}} \sum_{j \in I} \sum_k u_{ik} \otimes u_{kj} \right) = \sum_k u_{ik} \otimes (\xi_I)_k = x_i \otimes 1$$

Finally, for the group duals this quotient map is given by $\widehat{\Gamma}'_I \rightarrow \widehat{\Gamma}_I$, where $\Gamma'_I \subset \Gamma$ is the normal closure of Γ_I , and so this map can be indeed proper. See [1]. \square

2. ALGEBRAIC MANIFOLDS

We discuss in what follows the axiomatization of the affine homogeneous spaces, as algebraic submanifolds of the free sphere $S_{\mathbb{C},+}^{N-1}$. We use the following formalism:

Definition 2.1. A closed subset $X \subset S_{\mathbb{C},+}^{N-1}$ is called algebraic when

$$C(X) = C(S_{\mathbb{C},+}^{N-1}) / \left\langle P_i(x_1, \dots, x_N) = 0, \forall i \in I \right\rangle$$

for a certain family of noncommutative $*$ -polynomials $P_i \in \mathbb{C} \langle x_1, \dots, x_N \rangle$.

There are many examples of such manifolds, as for instance all the compact matrix quantum groups. Indeed, assuming that we have a closed subgroup $G \subset U_N^+$, by rescaling the standard coordinates we obtain an embedding $G \subset U_N^+ \subset S_{\mathbb{C},+}^{N^2-1}$, and the following result, coming from [14], shows that we have indeed an algebraic manifold:

Proposition 2.2. Given a closed subgroup $G \subset U_N^+$, with the corresponding fundamental corepresentations denoted $u \rightarrow v$, we have the formula

$$C(G) = C(U_N^+) / \left(T \in \text{Hom}(u^{\otimes k}, u^{\otimes l}), \forall k, l, \forall T \in \text{Hom}(v^{\otimes k}, v^{\otimes l}) \right)$$

with $k, l = \dots \circ \bullet \bullet \circ \bullet \dots$ being colored integers, with the tensor power conventions $w^\circ = w, w^\bullet = \bar{w}, w^{xy} = w^x \otimes w^y$, and with the notation $\text{Hom}(r, p) = \{T \mid Tr = pT\}$.

Proof. For any choice of two colored integers k, l and of an intertwiner $T \in \text{Hom}(v^{\otimes k}, v^{\otimes l})$, the formula $T \in \text{Hom}(u^{\otimes k}, u^{\otimes l})$, with $u = (u_{ij})$ being the fundamental corepresentation of $C(U_N^+)$, corresponds to a collection of N^{k+l} relations between the variables u_{ij} . By dividing now $C(U_N^+)$ by the ideal generated by all these relations, when k, l and T vary, we obtain a certain algebra A , which is the algebra on the right in the statement.

It is clear that we have a surjective morphism $A \rightarrow C(G)$, and by using Woronowicz's Tannakian results in [14], this surjective morphism follows to be an isomorphism. For a short, recent proof of this fact, using basic Hopf algebra theory, see [9]. \square

In relation now with the affine homogeneous spaces, we have:

Proposition 2.3. Any affine homogeneous space $X_{G,I} \subset S_{\mathbb{C},+}^{N-1}$ is algebraic, with

$$\sum_{i_1 \dots i_k} \xi_{i_1 \dots i_k} x_{i_1}^{e_1} \dots x_{i_k}^{e_k} = \frac{1}{\sqrt{|I|^k}} \sum_{b_1 \dots b_k \in I} \xi_{b_1 \dots b_k} \quad \forall k, \forall \xi \in \text{Fix}(u^{\otimes k})$$

as defining relations. Moreover, we can use vectors ξ belonging to a basis of $\text{Fix}(u^{\otimes k})$.

Proof. This follows indeed from the various results in section 1, and more specifically from Proposition 1.6, by using the identifications made in Theorem 1.7. \square

In order to reach to a more categorical description of $X_{G,I}$, the idea will be that of using Frobenius duality. We use colored indices, and we denote by $k \rightarrow \bar{k}$ the operation on the colored indices which consists in reversing the index, and switching all the colors. Also, we agree to identify the linear maps $T : (\mathbb{C}^N)^{\otimes k} \rightarrow (\mathbb{C}^N)^{\otimes l}$ with the corresponding rectangular matrices $T \in M_{N^l \times N^k}(\mathbb{C})$, written $T = (T_{i_1 \dots i_l, j_1 \dots j_k})$. With these conventions, the precise formulation of Frobenius duality that we will need is as follows:

Proposition 2.4. *We have an isomorphism of complex vector spaces*

$$T \in \text{Hom}(u^{\otimes k}, u^{\otimes l}) \longleftrightarrow \xi \in \text{Fix}(u^{\otimes l} \otimes u^{\otimes \bar{k}})$$

given by the formulae $T_{i_1 \dots i_l, j_1 \dots j_k} = \xi_{i_1 \dots i_l, j_k \dots j_1}$ and $\xi_{i_i \dots i_l, j_1 \dots j_k} = T_{i_1 \dots i_l, j_k \dots j_1}$.

Proof. This is a well-known result, which follows from the general theory in [13]. To be more precise, given integers $K, L \in \mathbb{N}$, consider the following standard isomorphism, which in matrix notation makes $T = (T_{IJ}) \in M_{L \times K}(\mathbb{C})$ correspond to $\xi = (\xi_{IJ})$:

$$T \in \mathcal{L}(\mathbb{C}^{\otimes K}, \mathbb{C}^{\otimes L}) \longleftrightarrow \xi \in \mathbb{C}^{\otimes L+K}$$

Given now two arbitrary corepresentations $v \in M_K(C(G))$ and $w \in M_L(C(G))$, the abstract Frobenius duality result established in [13] states that the above isomorphism restricts into an isomorphism of vector spaces, as follows:

$$T \in \text{Hom}(v, w) \longleftrightarrow \xi \in \text{Fix}(w \otimes \bar{v})$$

In our case, we can apply this result with $v = u^{\otimes k}$ and $w = u^{\otimes l}$. Since, according to our conventions, we have $\bar{v} = u^{\otimes \bar{k}}$, this gives the isomorphism in the statement. \square

With the above result in hand, we can enhance the construction of $X_{G,I}$, as follows:

Theorem 2.5. *Any affine homogeneous space $X_{G,I}$ is algebraic, with*

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} (x_{j_1}^{f_1} \dots x_{j_k}^{f_k})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

for any k, l , and any $T \in \text{Hom}(u^{\otimes k}, u^{\otimes l})$, as defining relations.

Proof. We must prove that the relations in the statement are satisfied, over $X_{G,I}$. We know from Proposition 2.3 above, with $k \rightarrow l\bar{k}$, that the following relation holds:

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} \xi_{i_1 \dots i_l, j_k \dots j_1} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} x_{j_k}^{\bar{f}_k} \dots x_{j_1}^{\bar{f}_1} = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} \xi_{b_1 \dots b_l, c_k \dots c_1}$$

In terms of the matrix $T_{i_1 \dots i_l, j_1 \dots j_k} = \xi_{i_1 \dots i_l, j_k \dots j_1}$ from Proposition 2.3, we obtain:

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} x_{j_k}^{\bar{f}_k} \dots x_{j_1}^{\bar{f}_1} = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

But this gives the formula in the statement, and we are done. \square

3. TANNAKIAN DUALITY

In this section we state and prove our main result. The description of the affine homogeneous spaces found in Theorem 2.5 above suggests the following notion:

Definition 3.1. Given an algebraic submanifold $X \subset S_{\mathbb{C},+}^{N-1}$ and a subset $I \subset \{1, \dots, N\}$, we say that X is I -affine when $C(X)$ is presented by relations of type

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} (x_{j_1}^{f_1} \dots x_{j_k}^{f_k})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

with the operators T belonging to certain linear spaces $F(k, l) \subset M_{N^l \times N^k}(\mathbb{C})$, which altogether form a tensor category $F = (F(k, l))$.

According to Theorem 2.5, any affine homogeneous space $X_{G,I}$ is an I -affine manifold, with the corresponding tensor category being the one associated to the quantum group $G \subset U_N^+$ which produces it, formed by the linear spaces $F(k, l) = \text{Hom}(u^{\otimes k}, u^{\otimes l})$.

We will need some basic facts regarding the quantum affine actions. Following Definition 1.2, we say that a closed subgroup $G \subset U_N^+$ acts affinely on a closed subset $X \subset S_{\mathbb{C},+}^{N-1}$ when the formula $\Phi(x_i) = \sum_j u_{ij} \otimes x_j$ defines a morphism of C^* -algebras.

We have the following standard result, from [4], inspired from [7], [8]:

Proposition 3.2. Given an algebraic manifold $X \subset S_{\mathbb{C},+}^{N-1}$, the quantum group

$$G^+(X) = \max \left\{ G \subset U_N^+ \mid G \curvearrowright X \right\}$$

exists, and is unique. We call it affine quantum isometry group of X .

Proof. In order to have a universal coaction, the relations defining $G^+(X) \subset U_N^+$ must be those making $x_i \rightarrow X_i = \sum_j u_{ij} \otimes x_j$ a morphism of algebras. Thus, in order to construct $G^+(X)$, we just have to clarify how the relations $P_\alpha(x_1, \dots, x_N) = 0$ defining X are interpreted inside $C(U_N^+)$. So, pick one such polynomial, $P = P_\alpha$, and write it:

$$P(x_1, \dots, x_N) = \sum_r \alpha_r \cdot x_{i_1}^{r_1} \dots x_{i_{s(r)}}^{r_{s(r)}}$$

Now if we formally replace each coordinate $x_i \in C(X)$ by the corresponding element $X_i = \sum_j u_{ij} \otimes x_j \in C(U_N^+) \otimes C(X)$, the following formula must hold:

$$P(X_1, \dots, X_N) = \sum_r \alpha_r \sum_{j_1^r \dots j_{s(r)}^r} u_{i_1^r j_1^r} \dots u_{i_{s(r)}^r j_{s(r)}^r} \otimes x_{j_1^r} \dots x_{j_{s(r)}^r}$$

Thus the relations $P(X_1, \dots, X_N) = 0$ correspond indeed to certain polynomial relations between the standard generators $u_{ij} \in C(U_N^+)$, and this gives the result. See [4]. \square

Now by getting back to our questions, let us study the quantum isometry groups of the manifolds $X \subset S_{\mathbb{C},+}^{N-1}$ which are I -affine. We have here the following result:

Proposition 3.3. For an I -affine manifold $X \subset S_{\mathbb{C},+}^{N-1}$ we have

$$G \subset G^+(X)$$

where $G \subset U_N^+$ is the Tannakian dual of the associated tensor category F .

Proof. We recall from the proof of Proposition 3.2 above that the relations defining $G^+(X)$ are those expressing the vanishing of the following quantities:

$$P(X_1, \dots, X_N) = \sum_r \alpha_r \sum_{j_1^r \dots j_{s(r)}^r} u_{i_1^r j_1^r} \dots u_{i_{s(r)}^r j_{s(r)}^r} \otimes x_{j_1^r} \dots x_{j_{s(r)}^r}$$

In the case of an I -affine manifold, the defining relations are those from Definition 3.1 above, with the corresponding polynomials P being indexed by the elements of F . But the vanishing of the associated relations $P(X_1, \dots, X_N) = 0$ corresponds precisely to the Tannakian relations defining $G \subset U_N^+$, and so we obtain $G \subset G^+(X)$, as claimed. \square

We have now all the needed ingredients, and we can prove:

Theorem 3.4. *Assuming that an algebraic manifold $X \subset S_{\mathbb{C},+}^{N-1}$ is I -affine, with associated tensor category F , the following happen:*

- (1) *We have an inclusion $G \subset G^+(X)$, where G is the Tannakian dual of F .*
- (2) *X is an affine homogeneous space, $X = X_{G,I}$, over this quantum group G .*

Proof. In the context of Definition 3.1, the tensor category F there gives rise, by the Tannakian result in Proposition 2.2, to a quantum group $G \subset U_N^+$. What is left now is to construct the affine space morphisms α, Φ , and the proof here goes as follows:

- (1) Construction of α . We want to construct a morphism, as follows:

$$\alpha : C(X) \rightarrow C(G) \quad : \quad x_i \rightarrow X_i = \frac{1}{\sqrt{|I|}} \sum_{j \in I} u_{ij}$$

In view of Definition 3.1, we must therefore prove that we have:

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} X_{i_1}^{e_1} \dots X_{i_l}^{e_l} (X_{j_1}^{f_1} \dots X_{j_k}^{f_k})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

By replacing the variables X_i by their above values, we want to prove that:

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} \sum_{r_1 \dots r_l \in I} \sum_{s_1 \dots s_k \in I} T_{i_1 \dots i_l, j_1 \dots j_k} u_{i_1 r_1}^{e_1} \dots u_{i_l r_l}^{e_l} (u_{j_1 s_1}^{f_1} \dots u_{j_k s_k}^{f_k})^* = \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

Now observe that from the relation $T \in \text{Hom}(u^{\otimes k}, u^{\otimes l})$ we obtain:

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} u_{i_1 r_1}^{e_1} \dots u_{i_l r_l}^{e_l} (u_{j_1 s_1}^{f_1} \dots u_{j_k s_k}^{f_k})^* = T_{r_1 \dots r_l, s_1 \dots s_k}$$

Thus, by summing over indices $r_i \in I$ and $s_i \in I$, we obtain the desired formula.

- (2) Construction of Φ . We want to construct a morphism, as follows:

$$\Phi : C(X) \rightarrow C(G) \otimes C(X) \quad : \quad x_i \rightarrow X_i = \sum_j u_{ij} \otimes x_j$$

But this is precisely the coaction map constructed in Proposition 3.3 above.

(3) Proof of the ergodicity. If we go back here to Proposition 1.4, we see that the ergodicity condition is equivalent to a number of Tannakian conditions, which are automatic in our case. Thus, the ergodicity condition is automatic, and we are done. \square

4. FURTHER RESULTS

The Tannakian result obtained in section 3 above, based on the notion of I -affine manifold from Definition 3.1, remains quite theoretical. The problem is that Definition 3.1 still makes reference to a tensor category, and so the abstract characterization of the affine homogeneous spaces that we obtain in this way is not totally intrinsic.

We believe that some deeper results should hold as well. To be more precise, the work on noncommutative spheres in [4] suggests that the relevant category F should appear in a more direct way from X . In analogy with Definition 3.1, let us formulate:

Definition 4.1. *Given a submanifold $X \subset S_{\mathbb{C},+}^{N-1}$ and a subset $I \subset \{1, \dots, N\}$, we let $F_{X,I}(k, l) \subset M_{N^l \times N^k}(\mathbb{C})$ be the linear space of linear maps T such that*

$$\sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} (x_{j_1}^{f_1} \dots x_{j_k}^{f_k})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k}$$

holds over X . We say that X is I -saturated when $F_{X,I} = (F_{X,I}(k, l))$ is a tensor category, and the collection of the above relations presents the algebra $C(X)$.

Observe that any I -saturated manifold is automatically I -affine. The point is that the results in [4] seem to suggest that the converse of this fact should hold, in the sense that any I -affine manifold should be automatically I -saturated. Such a result would of course substantially improve Theorem 3.4 above, and make it ready for applications.

We do not have a proof of this fact, but we would like to present now a few preliminary observations on this subject. First of all, we have the following result:

Proposition 4.2. *The linear spaces $F_{X,I}(k, l) \subset M_{N^l \times N^k}(\mathbb{C})$ constructed above have the following properties:*

- (1) *They contain the units.*
- (2) *They are stable by conjugation.*
- (3) *They satisfy the Frobenius duality condition.*

Proof. All these assertions are elementary, as follows:

- (1) Consider indeed the unit map. The associated relation is:

$$\sum_{i_1 \dots i_k} x_{i_1}^{e_1} \dots x_{i_k}^{e_k} (x_{i_1}^{e_1} \dots x_{i_k}^{e_k})^* = 1$$

But this relation holds indeed, due to the defining relations for $S_{\mathbb{C},+}^{N-1}$.

(2) We have indeed the following sequence of equivalences:

$$\begin{aligned}
& T^* \in F_{X,I}(l, k) \\
\iff & \sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{j_1 \dots j_k, i_1 \dots i_l}^* x_{j_1}^{f_1} \dots x_{j_k}^{f_k} (x_{i_1}^{e_1} \dots x_{i_l}^{e_l})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{c_1 \dots c_k, b_1 \dots b_l}^* \\
\iff & \sum_{i_1 \dots i_l} \sum_{j_1 \dots j_k} T_{i_1 \dots i_l, j_1 \dots j_k} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} (x_{j_1}^{f_1} \dots x_{j_k}^{f_k})^* = \frac{1}{\sqrt{|I|^{k+l}}} \sum_{b_1 \dots b_l \in I} \sum_{c_1 \dots c_k \in I} T_{b_1 \dots b_l, c_1 \dots c_k} \\
\iff & T \in F_{X,I}(k, l)
\end{aligned}$$

(3) We have indeed a correspondence $T \in F_{X,I}(k, l) \longleftrightarrow \xi \in F_{X,I}(\emptyset, \bar{l}k)$, given by the usual formulae for the Frobenius isomorphism, from Proposition 2.4. \square

Based on the above result, we can now formulate our observations, as follows:

Theorem 4.3. *Given a closed subgroup $G \subset U_N^+$, and an index set $I \subset \{1, \dots, N\}$, consider the corresponding affine homogeneous space $X_{G,I} \subset S_{\mathbb{C},+}^{N-1}$.*

- (1) $X_{G,I}$ is I -saturated precisely when the collection of spaces $F_{X,I} = (F_{X,I}(k, l))$ is stable under compositions, and under tensor products.
- (2) We have $F_{X,I} = F$ precisely when $\sum_{j_1 \dots j_l \in I} (\sum_{i_1 \dots i_l} \xi_{i_1 \dots i_l} u_{i_1 j_1}^{e_1} \dots u_{i_l j_l}^{e_l} - \xi_{j_1 \dots j_l}) = 0$ implies $\sum_{i_1 \dots i_l} \xi_{i_1 \dots i_l} u_{i_1 j_1}^{e_1} \dots u_{i_l j_l}^{e_l} - \xi_{j_1 \dots j_l} = 0$, for any j_1, \dots, j_l .

Proof. We use the fact, coming from Theorem 2.5, that with $F(k, l) = \text{Hom}(u^{\otimes k}, u^{\otimes l})$, we have inclusions of vector spaces $F(k, l) \subset F_{X,I}(k, l)$. Moreover, once again by Theorem 2.5, the relations coming from the elements of the category formed by the spaces $F(k, l)$ present $X_{G,I}$. Thus, the relations coming from the elements of $F_{X,I}$ present $X_{G,I}$ as well.

With this observation in hand, our assertions follow from Proposition 4.2:

(1) According to Proposition 4.2 (1) and (2) the unit and conjugation axioms are satisfied, so the spaces $F_{X,I}(k, l)$ form a tensor category precisely when the remaining axioms, namely the composition and the tensor product one, are satisfied. Now by assuming that these two axioms are satisfied, X follows to be I -saturated, by the above observation.

(2) Since we already have inclusions in one sense, the equality $F_{X,I} = F$ from the statement means that we must have inclusions in the other sense, as follows:

$$F_{X,I}(k, l) \subset F(k, l)$$

By using now Proposition 4.2 (3), it is enough to discuss the case $k = 0$. And here, assuming that we have $\xi \in F_{X,I}(0, l)$, the following condition must be satisfied:

$$\sum_{i_1 \dots i_l} \xi_{i_1 \dots i_l} x_{i_1}^{e_1} \dots x_{i_l}^{e_l} = \sum_{j_1 \dots j_l \in I} \xi_{j_1 \dots j_l}$$

By applying now the morphism $\alpha : C(X_{G,I}) \rightarrow C(G)$, we deduce that we have:

$$\sum_{i_1 \dots i_l} \xi_{i_1 \dots i_l} \sum_{j_1 \dots j_l \in I} u_{i_1 j_1}^{e_1} \dots u_{i_l j_l}^{e_l} = \sum_{j_1 \dots j_l \in I} \xi_{j_1 \dots j_l}$$

Now recall that $F(0, l) = \text{Fix}(u^{\otimes l})$ consists of the vectors ξ satisfying:

$$\sum_{i_1 \dots i_l} \xi_{i_1 \dots i_l} u_{i_1 j_1}^{e_1} \dots u_{i_l j_l}^{e_l} = \xi_{j_1 \dots j_l}, \forall j_1, \dots, j_l$$

We are therefore led to the conclusion in the statement. \square

It is quite unclear on how to advance on these questions, and a more advanced algebraic trick, in the spirit of those used in [4], seems to be needed. Nor is it clear on how to explicitly “capture” the relevant subgroup $G \subset G^+(X)$, in terms of our given manifold $X = X_{G,I}$, in a direct, geometric way. Summarizing, further improving Theorem 3.4 above is an interesting question, that we would like to raise here.

REFERENCES

- [1] T. Banica, Weingarten integration over noncommutative homogeneous spaces, *Ann. Math. Blaise Pascal* **24** (2017), 195–224.
- [2] T. Banica and B. Collins, Integration over compact quantum groups, *Publ. Res. Inst. Math. Sci.* **43** (2007), 277–302.
- [3] T. Banica and D. Goswami, Quantum isometries and noncommutative spheres, *Comm. Math. Phys.* **298** (2010), 343–356.
- [4] T. Banica and S. Mészáros, Uniqueness results for noncommutative spheres and projective spaces, *Illinois J. Math.* **59** (2015), 219–233.
- [5] B. Collins and S. Matsumoto, Weingarten calculus via orthogonality relations: new applications, preprint 2016.
- [6] B. Collins and P. Śniady, Integration with respect to the Haar measure on the unitary, orthogonal and symplectic group, *Comm. Math. Phys.* **264** (2006), 773–795.
- [7] D. Goswami, Quantum group of isometries in classical and noncommutative geometry, *Comm. Math. Phys.* **285** (2009), 141–160.
- [8] D. Goswami, Existence and examples of quantum isometry groups for a class of compact metric spaces, *Adv. Math.* **280** (2015), 340–359.
- [9] S. Malacarne, Woronowicz’s Tannaka-Krein duality and free orthogonal quantum groups, preprint 2016.
- [10] S. Raum, Isomorphisms and fusion rules of orthogonal free quantum groups and their complexifications, *Proc. Amer. Math. Soc.* **140** (2012), 3207–3218.
- [11] S. Wang, Free products of compact quantum groups, *Comm. Math. Phys.* **167** (1995), 671–692.
- [12] D. Weingarten, Asymptotic behavior of group integrals in the limit of infinite rank, *J. Math. Phys.* **19** (1978), 999–1001.
- [13] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [14] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35–76.

T.B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CERGY-PONTOISE, F-95000 CERGY-PONTOISE, FRANCE. teo.banica@gmail.com