

QUANTUM PERMUTATIONS, HADAMARD MATRICES, AND THE SEARCH FOR MATRIX MODELS

TEODOR BANICA

ABSTRACT. This is a presentation of recent work on quantum permutation groups, complex Hadamard matrices, and the connections between them. A long list of problems is included. We include as well some conjectural statements, about matrix models.

Dedicated to Professor S.L. Woronowicz, on the occasion of his 70-th birthday.

CONTENTS

Introduction	1
Acknowledgements	2
1. Quantum permutations	3
2. Representation theory	7
3. Actions on finite spaces	12
4. Hadamard matrices	18
5. The correspondence	24
6. Matrix models	28
Conclusion	31
References	31

INTRODUCTION

A complex Hadamard matrix is a square matrix $H \in M_n(\mathbb{C})$ whose entries are on the unit circle, $|H_{ij}| = 1$, and whose rows are pairwise orthogonal, when regarded as elements of \mathbb{C}^n . It follows from definitions that the columns are pairwise orthogonal as well.

The basic example is the Fourier matrix $F_n = (w^{ij})$, where $w = e^{2\pi i/n}$. The name comes from the fact that F_n is the matrix of the discrete Fourier transform, over \mathbb{Z}_n . This is in fact the one and only basic example, typically the only known one when n is prime.

However, lots of other interesting Hadamard matrices, each one having its own story, do exist. For instance in the real case the Hadamard conjecture, going back to the 19th century, states that there is a such a matrix $H \in M_n(\mathbb{R})$, for any $n \in 4\mathbb{N}$.

2000 *Mathematics Subject Classification.* 46L65 (05B20).

Key words and phrases. Quantum permutation, Hadamard matrix.

Due to their remarkable combinatorial properties, the complex Hadamard matrices appear in a wealth of concrete situations, in connection with subfactors, spin models, knot invariants, planar algebras, quantum groups, and various aspects of combinatorics, functional analysis, representation theory, and quantum physics.

The present article is a survey on complex Hadamard matrices, quantum permutation groups, and the connections between them. The idea is very simple:

- (1) First, the Fourier matrix F_n is by definition cyclic, and is in fact naturally associated to \mathbb{Z}_n . This suggests that any complex Hadamard matrix H should be in fact associated to a group-like object G , describing its symmetries.
- (2) A glimpse at the various complex Hadamard matrices (different from F_n) which have been constructed leads to the quick conclusion that their symmetries are rather of “alien” nature, and cannot be described by a usual group G .
- (3) So, we should rather look for a quantum group. And, with this idea in mind, the solution appears: the orthogonality condition on the rows of H reminds the “magic” condition on the coordinates of the quantum permutation group S_n^+ .
- (4) With a bit more work, one can associate to any complex Hadamard matrix $H \in M_n(\mathbb{C})$ a certain quantum permutation group $G \subset S_n^+$. As a first example, associated in this way to the Fourier matrix F_n is the usual cyclic group \mathbb{Z}_n .

There’s perhaps at this point some need for explanations: the complex Hadamard matrices are such concrete and beautiful objects, so why doing all this?

The problem is that any complex Hadamard matrix H produces a certain sequence of numbers $c_k \in \mathbb{N}$, which are extremely hard to compute. These numbers, called “quantum invariants”, appeared from the work of Popa and Jones on subfactor theory, and play a key role in most of the quantum algebraic contexts where H appears. Their formal definition is $c_k = \dim(P_k)$, where $P = (P_k)$ is the planar algebra associated to H .

The point now is that, with the above approach, the numbers c_k have a very simple interpretation. Indeed, they appear as moments of the main character of G :

$$c_k = \int_G \text{Tr}(g)^k dg$$

This formula is of course quite encouraging. For instance it shows that, from the point of view of quantum algebra in a very large sense, G is indeed the correct symmetry group of H . Of course, these results are quite new, and the correspondence $H \rightarrow G$ still needs to be developed in order to reach to some concrete results. It is our hope that so will be the case, and in the present article we explain the state-of-art of the subject.

The paper is organized as follows: in 1-3 we discuss the quantum permutation groups, in 4 we discuss the complex Hadamard matrices, in 5 we describe the correspondence and its basic properties, and in 6 we present some random matrix speculations.

Acknowledgements. It would be impossible to make a full list of those who have made this theory possible, but at the beginning we certainly have Professor S.L. Woronowicz, to

whom this article is dedicated. It is also a pleasure to thank Julien Bichon, Benoît Collins and Jean-Marc Schlenker for their key contribution to the subject, Stephen Curran, Sonia Natale, Adam Skalski and Roland Speicher for their work on quantum permutations, and Dietmar Bisch, Vaughan Jones and Remus Nicoara for various subfactor discussions.

The present article is based on lecture notes from minicourses given at the University of Caen in Fall 2010, and at the Erwin Schrödinger Institute in Vienna in Spring 2011, and I am grateful to Roland Vergnioux and Dan Voiculescu for the invitations.

This work was supported by the ANR grants “Galosint” and “Granma”.

1. QUANTUM PERMUTATIONS

Consider the embedding $S_n \subset O_n$ given by the usual permutation matrices, $\sigma(e_j) = e_{\sigma(j)}$. The matrix entries of a permutation $\sigma \in S_n$ are then:

$$\sigma_{ij} = \begin{cases} 1 & \text{if } \sigma(j) = i \\ 0 & \text{if not} \end{cases}$$

Let us introduce now the functions $u_{ij} : S_n \rightarrow \mathbb{C}$ given by $u_{ij}(\sigma) = \sigma_{ij}$. These are the n^2 coordinates on S_n , viewed as an algebraic group.

We can see that the matrix $u = (u_{ij})$ is magic, in the following sense:

Definition 1.1. *A square matrix $u \in M_n(A)$ is called “magic” if all its entries are projections, summing up to 1 on each row and column.*

Here of course A is a C^* -algebra, and a projection is by definition an element $p \in A$ satisfying $p = p^* = p^2$. In the above example, the algebra is $A = C(S_n)$.

Consider now the multiplication map $(\sigma, \tau) \rightarrow \sigma\tau$. At the level of functions, this corresponds to a map $\Delta : A \rightarrow A \otimes A$ called comultiplication, given by $\Delta f(\sigma, \tau) = f(\sigma\tau)$. On the above elements u_{ij} , the action of Δ is extremely simple:

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}$$

In addition, we know from the Stone-Weierstrass theorem that the coordinate functions u_{ij} generate A . With a bit more work one can prove that these generators u_{ij} satisfy no other relations besides the magic ones, so we have reached to the following conclusion:

Theorem 1.2. *$C(S_n)$ is the universal commutative C^* -algebra generated by the entries of a $n \times n$ magic matrix u , and its comultiplication is given by $\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj}$.*

Summarizing, we have a fully satisfactory description of the Hopf algebra $C(S_n)$. Observe that the counit map $\varepsilon(f) = f(1)$ and the antipode map $Sf(\sigma) = f(\sigma^{-1})$ are given as well by very simple formulae on the standard generators, namely:

$$\varepsilon(u_{ij}) = \delta_{ij}, \quad S(u_{ij}) = u_{ji}$$

We recall now that a compact quantum group is an abstract object G , having no points in general, but which is described by a well-defined Hopf C^* -algebra $A = C(G)$.

The axioms for Hopf C^* -algebras, found by Woronowicz in [87], are as follows:

Definition 1.3. *A Hopf C^* -algebra is a C^* -algebra A , given with a morphism of C^* -algebras $\Delta : A \rightarrow A \otimes A$, called comultiplication, subject to the following conditions:*

- (1) *Coassociativity:* $(\Delta \otimes id)\Delta = (id \otimes \Delta)\Delta$.
- (2) *$\overline{\text{span}} \Delta(A)(A \otimes 1) = \overline{\text{span}} \Delta(A)(1 \otimes A) = A \otimes A$.*

The basic example is $A = C(G)$, where G is a compact group, with $\Delta f(g, h) = f(gh)$. The fact that Δ is coassociative corresponds to $(gh)k = g(hk)$, and the conditions in (2) correspond to the cancellation rules $gh = gk \implies h = k$ and $gh = kh \implies g = k$.

Conversely, any commutative Hopf C^* -algebra is of the form $C(G)$. Indeed, by the Gelfand theorem we have $A = C(G)$, with G compact space, and (1,2) above tell us that G is a semigroup with cancellation. By a well-known result, G follows to be a group.

The following key definition is due to Wang [84]:

Definition 1.4. *$C(S_n^+)$ is the universal C^* -algebra generated by the entries of a $n \times n$ magic matrix u , with comultiplication $\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj}$.*

Observe the similarity with Theorem 1.2. Wang proved that this algebra satisfies Woronowicz's axioms in Definition 1.3, so that S_n^+ is a compact quantum group.

The counit and antipode are given by the same formulae as in the classical case, namely $\varepsilon(u_{ij}) = \delta_{ij}$ and $S(u_{ij}) = u_{ji}$. Observe that the square of the antipode is the identity:

$$S^2 = id$$

As a first remark, we have a surjective morphism $C(S_n^+) \rightarrow C(S_n)$, which corresponds to a quantum group embedding $S_n \subset S_n^+$. The very first problem is whether this embedding is an isomorphism or not, and the answer here is as follows:

Proposition 1.5. *The quantum groups S_n^+ are as follows:*

- (1) *At $n = 1, 2, 3$ we have $S_n = S_n^+$.*
- (2) *At $n \geq 4$, S_n^+ is a non-classical, infinite compact quantum group.*

This is clear indeed at $n = 2$, where each magic matrix must be of the form:

$$u = \begin{pmatrix} p & 1-p \\ 1-p & p \end{pmatrix}$$

At $n = 3$ the statement is that any 9 projections forming a 3×3 magic matrix commute. This is not obvious, but can be seen with a Fourier over \mathbb{Z}_3 base change.

At $n = 4$ now, consider the following magic matrix:

$$u = \begin{pmatrix} p & 1-p & 0 & 0 \\ 1-p & p & 0 & 0 \\ 0 & 0 & q & 1-q \\ 0 & 0 & 1-q & q \end{pmatrix}$$

Here p, q can be any projections. If we choose $p, q \in B(H)$ to be free, then $\langle p, q \rangle$ is not commutative and infinite dimensional, and so must be $C(S_4^+)$.

Finally, at $n \geq 5$ the result follows from $S_k^+ \subset S_n^+$ for $k \leq n$.

The following result was proved as well by Wang in [84]:

Theorem 1.6. *$C(S_n^+)$ is the universal Hopf C^* -algebra coacting on \mathbb{C}^n . In other words, S_n^+ is the universal compact quantum group acting on $\{1, \dots, n\}$.*

The idea here is that a coaction map $\alpha(e_i) = \sum_j e_j \otimes u_{ji}$ is a morphism of algebras if and only if the matrix $u = (u_{ij})$ is magic. Thus, if we try to construct the universal algebra coacting on \mathbb{C}^n , we are naturally led to Definition 1.4. See Wang [84].

Let us go back now to the above proof of the fact that S_4^+ is infinite. If we denote by X the “quantum space” corresponding to the algebra $\langle p, q \rangle$, the above proof is:

$$X \subset S_4^+, |X| = \infty \implies |S_4^+| = \infty$$

The point now is that X is in fact a quantum group. Indeed, it is well-known that the algebra $\langle p, q \rangle$ generated by two free projections is always the same, regardless of the projections p, q that we choose. One model is provided by $p = (1+g)/2$ and $q = (1+h)/2$, where g, h are the standard generators of $D_\infty = \mathbb{Z}_2 * \mathbb{Z}_2$, and we get:

Proposition 1.7. *We have an embedding $\widehat{D}_\infty \subset S_4^+$.*

In order to explain the meaning of this statement, we have to go back to Definition 1.3. Recall that the basic example there was $A = C(G)$, with G compact group. The point is that there is another class of “basic examples”, namely the group algebras $A = C^*(\Gamma)$, with $\Delta(g) = g \otimes g$. So, the abstract dual $\widehat{\Gamma}$ of a discrete group Γ can be defined by:

$$C^*(\Gamma) = C(\widehat{\Gamma})$$

With this notion in hand, the above result simply tells us that the morphism $C(S_4^+) \rightarrow C^*(D_\infty)$ commutes with the comultiplications. But this is clear from definitions.

Proposition 1.7 raises several questions, the first one being the classification of group dual subgroups $\widehat{\Gamma} \subset S_n^+$. The answer here, found by Bichon in [37], is as follows:

Theorem 1.8. *For any quotient group $\mathbb{Z}_{n_1} * \dots * \mathbb{Z}_{n_k} \rightarrow \Gamma$ we have $\widehat{\Gamma} \subset S_n^+$, where $n = n_1 + \dots + n_k$. Any group dual subgroup of S_n^+ appears in this way.*

The proof of the first assertion is similar to that of Proposition 1.7, by using a diagonal concatenation of magic matrices. The proof of the converse is quite technical, see [37].

Recall now Cayley's theorem, stating that any finite quantum group G is a permutation group: $G \subset S_n$, with $n = |G|$. The above result, when applied to the quotient map $\mathbb{Z}_n^{*n} \rightarrow G$, shows that \widehat{G} is a quantum permutation group: $\widehat{G} \subset S_{n^2}^+$. So, the following problem appears: is any finite quantum group a quantum permutation group?

This question was recently answered in the negative [15]:

Theorem 1.9. *There exists a quantum group of finite order, namely $\mathbb{Z}_4 \# \widehat{S}_3$ having order 24, which is not a quantum permutation group.*

We refer to [15] for the proof, as well as for a number of positive results: for instance any quantum group whose order divides p^3 or pqr is a quantum permutation one.

The other question raised by Proposition 1.7 is that of understanding the structure of S_4^+ . The answer here, quite surprising, involves the following definition, from [9]:

Definition 1.10. *$C(SO_3^{-1})$ is the universal C^* -algebra generated by the entries of a 3×3 orthogonal matrix $a = (a_{ij})$, with the following relations:*

- (1) *Skew-commutation: $a_{ij}a_{kl} = \pm a_{kl}a_{ij}$, with sign $+$ if $i \neq k, j \neq l$, and $-$ if not.*
- (2) *Twisted determinant condition: $\sum_{\sigma \in S_3} a_{1\sigma(1)}a_{2\sigma(2)}a_{3\sigma(3)} = 1$.*

Consider indeed the matrix $a^+ = \text{diag}(1, a)$, corresponding to the action of SO_3^{-1} on \mathbb{C}^4 , and apply to it the Fourier transform over the Klein group $K = \mathbb{Z}_2 \times \mathbb{Z}_2$:

$$u = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a_{11} & a_{12} & a_{13} \\ 0 & a_{21} & a_{22} & a_{23} \\ 0 & a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \end{pmatrix}$$

The point is that this matrix is magic, and vice versa, i.e. the Fourier transform over K converts the relations in Definition 1.10 into the relations in Definition 1.1.

This gives the following result, proved in [9]:

Theorem 1.11. *We have $S_4^+ = SO_3^{-1}$, with the algebraic isomorphism given on the standard generators by the Fourier transform over the Klein group $K = \mathbb{Z}_2 \times \mathbb{Z}_2$.*

This isomorphism raises the problem of classifying the ‘‘quantum groups acting on 4 points’’, i.e. the subgroups of S_4^+ . The result here, obtained in [9], is as follows:

Theorem 1.12. *The following is the list of subgroups of $S_4^+ = SO_3^{-1}$:*

- (1) *Infinite quantum groups: $S_4^+, O_2^{-1}, \widehat{D}_\infty$.*
- (2) *Finite groups: S_4 , and its subgroups.*
- (3) *Finite group twists: S_4^{-1}, A_5^{-1} .*
- (4) *Series of twists: D_{2n}^{-1} ($n \geq 3$), DC_n^{-1} ($n \geq 2$).*
- (5) *A group dual series: \widehat{D}_n , with $n \geq 3$.*

A finer version of this result is provided by the following table, jointly classifying the subgroups of SO_3 and of S_4^+ , and containing as well the corresponding ADE graphs:

Subgroup of SO_3	ADE graph	Subgroup of S_4^+
$\mathbb{Z}_1, \mathbb{Z}_2, \mathbb{Z}_3, \mathbb{Z}_4$	$\tilde{A}_1, \tilde{A}_3, \tilde{A}_5, \tilde{A}_7$	$\mathbb{Z}_1, \mathbb{Z}_2, \mathbb{Z}_3, K$
$\mathbb{Z}_{2n-1}, \mathbb{Z}_{2n}$	$\tilde{A}_{4n-3}, \tilde{A}_{4n-1}$	\square, \widehat{D}_n
SO_2, O_2, SO_3	$A_{-\infty, \infty}, D_\infty, A_\infty$	$\widehat{D}_\infty, O_2^{-1}, S_4^+$
K, D_4	\tilde{D}_4, \tilde{D}_6	\mathbb{Z}_4, D_4
D_{2n-1}, D_{2n}	$\tilde{D}_{2n+1}, \tilde{D}_{2n+2}$	$\square, D_{2n}^{-1}/DC_n^{-1}$
\square, \square	\square, \square	D_1, S_3
A_4, S_4, A_5	$\tilde{E}_6, \tilde{E}_7, \tilde{E}_8$	$A_4, S_4/S_4^{-1}, A_5^{-1}$

Here the squares denote the various missing objects in the correspondence, and the slashes indicate the places where the correspondence is not one-to-one.

The meaning of the middle column is a bit formal, in the sense that the graphs there are Cayley type graphs for the corresponding groups or quantum groups. See [9].

2. REPRESENTATION THEORY

We have seen in the previous section that S_4^+ is understood quite well. In order to discuss now the general case, we need some basic notions of representation theory:

Definition 2.1. *A unitary representation of G is described by the corresponding corepresentation of $A = C(G)$, which is a unitary matrix $u \in M_n(A)$ satisfying:*

$$\begin{aligned} \Delta(u_{ij}) &= \sum_k u_{ik} \otimes u_{kj} \\ \varepsilon(u_{ij}) &= \delta_{ij} \\ S(u_{ij}) &= u_{ji}^* \end{aligned}$$

The representation is called “fundamental” if its coefficients generate A .

Woronowicz proved that the compact quantum groups are semisimple. More precisely, his results include a Peter-Weyl type theory [85], and a Tannaka-Krein duality [86].

The main problem is to classify the irreducible representations of G , and to find their fusion rules. In the self-adjoint case, $u = \bar{u}$, it is known that each $r \in Irr(G)$ appears in a certain tensor power $u^{\otimes k}$. So, the problem is to compute the spaces $End(u^{\otimes k})$.

It is convenient to enlarge the attention to the following family of spaces:

$$C(k, l) = Hom(u^{\otimes k}, u^{\otimes l})$$

The point is that these spaces form a tensor category, and Woronowicz’s Tannakian duality results in [86] can be effectively used for their computation.

Let $P(k, l)$ be the set of partitions between k upper points, and l lower points.

Definition 2.2. Associated to $\pi \in P(k, l)$ is the linear map $T_\pi : (\mathbb{C}^n)^{\otimes k} \rightarrow (\mathbb{C}^n)^{\otimes l}$,

$$T(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1 \dots j_l} \delta_\pi \begin{pmatrix} i_1 & \dots & i_k \\ j_1 & \dots & j_l \end{pmatrix} e_{j_1} \otimes \dots \otimes e_{j_l}$$

where $\delta_\pi = 1$ if the indices contained in any block of π are all equal, and $\delta_\pi = 0$ if not.

It follows from definitions that the spaces $\text{span}(T_\pi | \pi \in P(k, l))$ form a tensor category. Indeed, the composition and tensor product of linear maps correspond to the vertical and horizontal concatenation of the partitions, and the $*$ operation corresponds to the upside-down turning of partitions. Moreover, it is easy to see that for any $\pi \in P(k, l)$ we have $T_\pi \in \text{Hom}(u^{\otimes k}, u^{\otimes l})$, where u is the fundamental representation of S_n .

Let also $NC(k, l) \subset P(k, l)$ be the set of noncrossing partitions.

Theorem 2.3. The spaces $\text{Hom}(u^{\otimes k}, u^{\otimes l})$ for S_n, S_n^+ are as follows:

- (1) For S_n we obtain $\text{span}(T_\pi | \pi \in P(k, l))$.
- (2) For S_n^+ we obtain $\text{span}(T_\pi | \pi \in NC(k, l))$.

As already mentioned, the inclusion $\text{span}(T_\pi | \pi \in P(k, l)) \subset \text{Hom}(u^{\otimes k}, u^{\otimes l})$ is easy to check for S_n . The reverse inclusion can be justified by several methods, none of them being trivial. For instance this inclusion follows from the classical Tannaka duality.

In the free case now, we can use the embedding $S_n \subset S_n^+$, which at the level of Hom spaces gives reverse inclusions $\text{Hom}(u^{\otimes k}, u^{\otimes l}) \subset \text{span}(T_\pi | \pi \in P(k, l))$. Now since S_n^+ was obtained from S_n by “removing commutativity”, at the Tannakian level we should “remove the basic crossing X ”, and so we reach to the above sets $NC(k, l)$.

In this latter proof we have used of course the results in [86]. See [17].

We recall that the law of a self-adjoint variable $a \in (A, \varphi)$ is the real probability measure having as moments the numbers $\varphi(a^k)$. In what follows φ will be the Haar functional.

Theorem 2.4. The law of the character $\chi = \text{Tr}(u)$ is as follows:

- (1) For S_n with $n \rightarrow \infty$ we obtain the Poisson law, $p = \frac{1}{e} \sum_{r=0}^{\infty} \frac{\delta_r}{r!}$.
- (2) For S_n^+ with $n \geq 4$ we obtain the free Poisson law, $\pi = \frac{1}{2\pi} \sqrt{4x^{-1} - 1} dx$.

The first result is well-known, and has a very beautiful proof. Indeed, since the standard coordinates on S_n are the characteristic functions $u_{ij} = \chi(\sigma | \sigma(j) = i)$, we obtain:

$$\chi = \sum_{i=1}^n \chi(\sigma | \sigma(i) = i) = |\text{fix}(\cdot)|$$

Now by using the well-known fact that a random permutation has no fixed points with probability $1/e$, hence has r fixed points with probability $1/(r!e)$, we get, as claimed:

$$\mathbb{P}(\chi(\sigma) = r) = \mathbb{P}(|\text{fix}(\sigma)| = r) \simeq \frac{1}{e} \cdot \frac{1}{r!}$$

Another proof, which works as well for S_n^+ , is as follows. Let $D(k, l)$ be the sets of diagrams appearing in Theorem 2.3. By using basic Peter-Weyl theory, we have:

$$\int \chi^k = \dim(\text{Fix}(u^{\otimes k})) \simeq |D(0, k)|$$

Here the sign at right corresponds to our $n \rightarrow \infty$ and $n \geq 4$ assumptions, and comes from the fact that the maps T_π are not exactly linearly independent. See [17]. Now, with this formula in hand, we conclude that the moments of χ are the Bell, respectively the Catalan numbers, and this gives the result.

Here is a purely functional analytic application of Theorem 2.4:

Corollary 2.5. *For $n \geq 5$ the discrete quantum group \widehat{S}_n^+ is not amenable.*

Indeed, recall the Kesten amenability criterion, which tells us that a discrete group $\Gamma = \langle g_1, \dots, g_n \rangle$, chosen with $\{g_i\} = \{g_i^{-1}\}$, is amenable if and only if:

$$n \in \text{Spec}(g_1 + \dots + g_n)$$

The point is that the Kesten operator is nothing but the trace of $u = \text{diag}(g_1, \dots, g_n)$, which is the fundamental representation of $\widehat{\Gamma}$. Thus, the Kesten criterion reads:

$$n \in \text{Spec}(\text{Tr}(u))$$

With a bit more work, one can show that the same result holds for any discrete quantum group Γ . See [5]. Thus, the above result follows from Theorem 2.4 (2) and from:

$$\text{Supp}(\pi) = [0, 4]$$

There are many interesting questions arising from Corollary 2.5, and from the work of Brannan [42], Fima [53], Kyed [65] and Vaes-Vergnioux [79]. In all cases, the question is whether the techniques there apply to S_n^+ , at $n \geq 5$.

Here is another corollary, this time of purely algebraic nature:

Corollary 2.6. *For $n \geq 4$ the irreducible representations of S_n^+ can be labeled by \mathbb{N} , and satisfy the Clebsch-Gordan rules: $r_a \otimes r_b = r_{|a-b|} + r_{|a-b|+1} + \dots + r_{a+b}$.*

This result can be indeed deduced from Theorem 2.4. The idea is that, with the initial data $r_0 = 1$ and $r_1 = u - 1$, we can construct a family of virtual representations r_a satisfying the Clebsch-Gordan rules. The problem is to prove that these virtual representations r_a are true representations, and are irreducible. But this can be done by recurrence on a , by using the fact that the moments of χ are the Catalan numbers:

$$\dim(\text{Fix}(u^{\otimes k})) = \frac{1}{k+1} \binom{2k}{k}$$

This proof is of course not very enlightening. Probably the best argument is as follows: let us enlarge attention to the groups $S(X)$ and quantum groups $S^+(X)$, where X can be any finite noncommutative space. We have then the following result:

Theorem 2.7. *If u denotes the fundamental representation of $S^+(X)$ then*

$$\text{Hom}(u^{\otimes k}, u^{\otimes l}) \simeq TL_n(k, l)$$

where TL_n denotes the Temperley-Lieb algebra of index $n = |X|$.

As a first remark, for a usual space $X = \{1, \dots, n\}$, this is exactly Theorem 2.3 (2) above, modulo a “fattening” of the diagrams. In general, the proof is based on the fact that the multiplication and unit of the algebra $C(X)$ can be represented as:

$$m = |\cup| \quad u = \cap$$

Now since $S^+(X)$ is the universal quantum group acting on $C(X)$, at the Tannakian level we get that its associated category is simply given by:

$$\langle m, u \rangle = \langle |\cup|, \cap \rangle = TL_n$$

With this conceptual result in hand, let us go back now to Corollary 2.6, and try to find a better proof. The point is that for the space $M_2 = \text{Spec}(M_2(\mathbb{C}))$ we have:

$$S^+(M_2) = S(M_2) = SO_3$$

On the other hand, it is well-known that under the Jones index assumption $n \geq 4$, the fusion ring for the category TL_n doesn't depend on n . Thus, the fusion rules for S_n^+ with $n \geq 4$ are the same as those for SO_3 , as stated in Corollary 2.6 above.

Problem 2.8. *Is it true that the compact quantum groups having Clebsch-Gordan fusion rules are exactly the quantum groups $S^+(X)$, with $|X| \geq 4$?*

The general method for attacking this type of problem is via Kazhdan-Wenzl reconstruction [62]. In the type B case the result here is due to Tuba and Wenzl [78].

Observe that the above discussion shows that with $q < 0$ given by $n = (q + q^{-1})^2$ we have a monoidal equivalence [38], i.e. an equivalence of tensor categories, as follows:

$$S_n^+ \sim SO_3^q$$

Observe also that the dimensions of the representations of S_n^+ with $n \geq 5$ are “exotic”. More precisely, if x, y are the roots of $X^2 - (n-2)X + 1 = 0$, then:

$$\dim(r_a) = \frac{x^{a+1} - y^{a+1}}{x - y} + \frac{x^a - y^a}{x - y}$$

There are as well a few other algebraic questions, coming from the work of Collins-Härtel-Thom [43], De Commer [48], Sołtan [74], Vergnioux [80] and Voigt [82].

Let us go back now to Theorem 2.4. One problem is to compute the value of the Haar functional on any element of type $u_{i_1 j_1} \dots u_{i_k j_k}$. The method here, developed for classical groups by Collins and Śniady in [44], applies as well to S_n, S_n^+ , and we have:

Theorem 2.9. *We have the Weingarten integration formula*

$$\int u_{i_1 j_1} \cdots u_{i_k j_k} = \sum_{\pi, \sigma \in D(k)} \delta_\pi(i) \delta_\sigma(j) W_{kn}(\pi, \sigma)$$

where $D(k) = D(0, k)$, and where $W_{kn} = G_{kn}^{-1}$, with $G_{kn}(\pi, \sigma) = n^{|\pi \vee \sigma|}$.

The proof of this formula, found in [17], is based on Theorem 2.3. Consider indeed the $n^k \times n^k$ matrix P obtained by integrating the coefficients of $u^{\otimes k}$. That is:

$$P = \left(\int u_{i_1 j_1} \cdots u_{i_k j_k} \right)_{i_1 \dots i_k, j_1 \dots j_k}$$

We know by Peter-Weyl theory that P is the projection on $Fix(u^{\otimes k})$, and from Theorem 2.3 that $Fix(u^{\otimes k})$ is spanned by $D(k)$. The Gram matrix of $D(k)$ is:

$$\langle T_\pi, T_\sigma \rangle = \sum_{i_1 \dots i_k} \delta_\pi(i) \delta_\sigma(i) = n^{|\pi \vee \sigma|}$$

By combining these ingredients and doing some linear algebra, this gives the result. As a first consequence, we have the following refinement of Theorem 2.4:

Theorem 2.10. *The asymptotic laws of truncated characters $\chi_t = \sum_{i=1}^{[tn]}$ are:*

(1) For S_n : the Poisson law, $p_t = e^{-t} \sum_{r=0}^{\infty} \frac{t^r}{r!} \delta_r$.

(2) For S_n^+ : the free Poisson law, $\pi_t = \max(1-t, 0) \delta_0 + \frac{\sqrt{4t-(x-1-t)^2}}{2\pi x} dx$.

This result is important for two reasons. First, it requires $n \rightarrow \infty$ for both S_n, S_n^+ , hence fixes the $n \rightarrow \infty$ vs. $n \geq 4$ issue in Theorem 2.4. And second, it proves that the ‘‘liberation’’ operation $S_n \rightarrow S_n^+$ is compatible with the Bercovici-Pata bijection [32].

The proof uses the Weingarten formula. With $s = [tn]$, we get:

$$\int \chi_t^k = \sum_{\pi, \sigma \in D(k)} s^{|\pi \vee \sigma|} W_{kn}(\pi, \sigma) = Tr(G_{ks} W_{kn})$$

Now since with $n \rightarrow \infty$, the Gram matrix becomes equal to its diagonal, $G_{kn} = \Delta_{kn} + O(n^{-1})$, by inverting we get as well $W_{kn} = \Delta_{kn}^{-1} + O(n^{-1})$, and this gives:

$$\int \chi_t^k = Tr(\Delta_{ks} \Delta_{kn}^{-1}) + O(n^{-1}) = \sum_{\pi \in D(k)} t^{|\pi|} + O(n^{-1})$$

According now to the general theory in [68], this shows that the cumulants for S_n , as well as the free cumulants for S_n^+ , are simply t, t, t, \dots , and this gives the result.

Let P_r be the Chebycheff polynomials, $P_0 = 1, P_1 = n$ and $P_{r+1} = nP_r - P_{r-1}$, and:

$$a_k = \frac{1}{k+1} \binom{2k}{k}, \quad f_{kr} = \binom{2k}{k-r} - \binom{2k}{k-r-1}$$

With these notations, the we have the following results:

Theorem 2.11. *The determinant of G_{kn} is as follows:*

- (1) For S_n : $\det(G_{kn}) = \prod_{\pi \in P(k)} \frac{n!}{(n-|\pi|)!}$.
- (2) For S_n^+ : $\det(G_{kn}) = (\sqrt{n})^{a_k} \prod_{r=1}^k P_r(\sqrt{n})^{d_{kr}}$, with $d_{kr} = f_{kr} - f_{k+1,r}$.

The classical result follows by an upper triangularization procedure. The free result uses Di Francesco's "meander determinant" in [49]. See [22].

We have as well the following generalization of Theorem 2.4:

Theorem 2.12. *The asymptotic law of $\chi^{(r)} = \text{Tr}(u^r)$ is as follows:*

- (1) For S_n : we obtain a linear combination of independent Poisson laws.
- (2) For S_n^+ : $\chi^{(1)}$ is free Poisson, $\chi^{(2)}$ is semicircular, $\chi^{(r)}$ with $r \geq 3$ is circular.

The classical result is due to Diaconis and Shahshahani [50]. Observe that in terms of the eigenvalue list $\lambda_1 \leq \dots \leq \lambda_n$, the variable in the statement is:

$$\text{Tr}(u^r) = \lambda_1^r + \dots + \lambda_n^r$$

In the free case the methods in [50] cannot work. In addition, a surprise appears at $r = 3$, where the variable in question is not self-adjoint:

$$\text{Tr}(u^3) = \sum_{ijk} u_{ij}u_{jk}u_{ki} \neq \sum_{ijk} u_{ki}u_{jk}u_{ij} = \text{Tr}(u^3)^*$$

We refer to our paper [24] for the proof for S_n^+ , which applies as well to S_n .

Question 2.13. *Is there a notion of "eigenvalues" for the quantum group S_n^+ , which is compatible with the above results?*

In other words, we would like to have a free analogue of the n eigenvalue functions $\lambda_1 \leq \dots \leq \lambda_n$, making the formulae in Theorem 2.12 work. As observed in [24], such eigenvalue functions can be constructed for certain related quantum groups O_n^*, H_n^* .

The main conceptual result regarding S_n, S_n^+ concerns their action on random variables. An action of $S = (S_n)$ or $S^+ = (S_n^+)$ on a sequence of random variables x_1, x_2, \dots is a coaction on the generated algebra, leaving invariant the joint distribution.

Theorem 2.14. *A sequence of random variables x_1, x_2, \dots is invariant iff:*

- (1) S case: x_i are independent and i.d. with respect to the tail algebra.
- (2) S^+ case: x_i are free and i.d. with respect to the tail algebra.

Here the first result is the classical de Finetti theorem, and the second result is the free De Finetti theorem, due to Köstler-Speicher [64]. A unified proof was given in [25].

3. ACTIONS ON FINITE SPACES

We have seen in the previous section that the quantum groups S_n^+ are understood quite well. In this section we extend some of these results, to subgroups $G \subset S_n^+$.

Consider a finite graph X , with edges colored by real numbers. The basic examples are the finite metric spaces, and the usual (monocolored) graphs. If we label the vertices $1, \dots, n$, the adjacency matrix becomes a real symmetric matrix, $d \in M_n(\mathbb{R})$.

Definition 3.1. $G^+(X)$ be the biggest quantum group acting on X , in the sense that $C(G^+(X))$ is the quotient of $C(S_n^+)$ by the relations coming from $du = ud$.

As a first remark, the classical version of $G^+(X)$ is the usual symmetry group $G(X)$. Indeed, with $u_{ij} = \chi(\sigma|_{\sigma(j)=i})$, the relation $du = ud$ reads:

$$d_{ij} = d_{\sigma(i)\sigma(j)}$$

Let us look now more carefully at the relation $du = ud$. Since d is a real symmetric matrix, we can consider its spectral decomposition:

$$d = \sum_{\lambda} \lambda \cdot P_{\lambda}$$

Also, for any $r \in \mathbb{R}$ we can consider the matrix d^r given by $d_{ij}^r = 1$ if $d_{ij} = r$, and $d_{ij}^r = 0$ if not. We obtain in this way the ‘‘color decomposition’’ of d :

$$d = \sum_r r \cdot d^r$$

Here, as in the case of the spectral decomposition, the sum is over nonzero terms.

Proposition 3.2. For a quantum group $G \subset S_n^+$, the following are equivalent:

- (1) We have $du = ud$, i.e. $G \subset G^+(X)$.
- (2) We have $P_{\lambda}u = uP_{\lambda}$, for any $\lambda \in \mathbb{R}$.
- (3) We have $d^r u = u d^r$, for any $r \in \mathbb{R}$.

Here the equivalence between (1) and (2) follows from definitions, and the equivalence with (3) follows by doing a bit of matrix analysis. These results are very useful, because they can be further combined: for instance u must commute with any color component P_{λ}^r of any spectral projection P_{λ} , and so on. This method of ‘‘spectral-color’’ decomposition proves to be quite efficient for the computation of $G^+(X)$. See [6].

Observe that the equivalence between (1) and (3) can be reformulated as follows, where X^r denotes the usual (monocolored) graph having adjacency matrix d^r :

$$G^+(X) = \bigcap_r G^+(X^r)$$

More generally now, let X be a finite noncommutative set, endowed with a self-adjoint operator $d : L^2(X) \rightarrow L^2(X)$, where the L^2 space is with respect the counting measure.

Definition 3.3. $G^+(X)$ be the biggest quantum group acting on X , in the sense that $C(G^+(X))$ is the quotient of $C(S^+(X_{\text{set}}))$ by the relations coming from $du = ud$.

Here we have used the quantum groups S^+ appearing in Theorem 2.7 above. This definition is of course very general, and covers for instance all the 0-dimensional spectral triples in the sense of Connes [45]. We will be back to these examples, a bit later.

Let $*$ be the free product of discrete quantum groups [83]:

$$C^*(\Gamma * \Lambda) = C^*(\Gamma) * C^*(\Lambda)$$

We denote by $\hat{*}$ the dual operation, at the level of compact quantum groups:

$$C(G \hat{*} H) = C(G) * C(H)$$

Finally, let us denote by $X_1 \sqcup \dots \sqcup X_k$ a disconnected union of graphs.

Proposition 3.4. *We have $G^+(X_1) \hat{*} \dots \hat{*} G^+(X_k) \subset G^+(X_1 \sqcup \dots \sqcup X_k)$.*

This follows from definitions, and might seem to be yet another triviality. However, let us work out the simplest example. For the graph formed by two segments we get:

$$\mathbb{Z}_2 \hat{*} \mathbb{Z}_2 \subset G^+(\text{---})$$

The dual free product on the left is by definition the dual of $\mathbb{Z}_2 * \mathbb{Z}_2 = D_\infty$. As for the quantum group of the right, we prefer to view it as the quantum symmetry group of the complement of $X = \text{---}$, which is $X^c = \square$. Thus, the above inclusion reads:

$$\widehat{D}_\infty \subset G^+(\square)$$

And this reminds of course the inclusion $\widehat{D}_\infty \subset S_4^+$ from Proposition 1.7 above, which was a bit the start of everything! More precisely, we have reached now to the following quite conceptual conclusion: “ S_4^+ is infinite simply because \widehat{D}_∞ acts on the square”.

Consider now an arbitrary union $kX = X \sqcup \dots \sqcup X$ (k terms).

Proposition 3.5. *If X is connected then $G^+(kX) = G^+(X) \wr_* S_k^+$.*

Here we use the free wreath product operation, constructed by Bichon in [36]:

$$C(G \wr_* H) = (C(G)^{*k} * C(H)) / \langle [u_{ij}^{(a)}, v_{ab}] = 0 \rangle$$

More generally now, let X, Y be finite graphs, and consider the graphs $X * Y$, $X \circ Y$ and $X \times Y$ having as vertex set $X_{set} \times Y_{set}$, and with the following edges:

$$(i, \alpha) \sim (j, \beta) \iff \begin{cases} \alpha \sim \beta \text{ or } \alpha = \beta, i \sim j & (* \text{ case}) \\ i \sim j, \alpha \sim \beta & (\circ \text{ case}) \\ i = j, \alpha \sim \beta \text{ or } i \sim j, \alpha = \beta & (\times \text{ case}) \end{cases}$$

In terms of the adjacency matrix the formulae are as follows, where \mathbb{I} denotes the square matrix filled with 1's:

$$d = \begin{cases} d_X \otimes 1 + \mathbb{I} \otimes d_Y & (* \text{ case}) \\ d_X \otimes d_Y & (\circ \text{ case}) \\ d_X \otimes 1 + 1 \otimes d_Y & (\times \text{ case}) \end{cases}$$

Let us first look at $*$. This product is obtained by putting a copy of X at each vertex of Y , so in particular we have $kX = X * Y_k$, where Y_k is the k -point graph.

Theorem 3.6. *Let X, Y be regular graphs, with spectra $\{\lambda_i\}, \{\mu_k\}$. If X is connected and*

$$\{\lambda_1 - \lambda_i | i \neq 1\} \cap \{-n\mu_k\} = \emptyset$$

*where n and λ_1 are the order and valence of X , then $G^+(X * Y) = G^+(X) \wr G^+(Y)$.*

This result, generalizing Proposition 3.5, was proved in [8]. Now let us look at \circ, \times . Under suitable assumptions on the eigenvalues of X, Y , we will get the spectral decomposition of d in terms of that of d_X, d_Y . If we assume in addition that X, Y are regular and connected, the 1-dimensional eigenspaces coming from the valence can be used for “splitting” the action of G^+ , and we have the following result, once again from [8]:

Theorem 3.7. *Let X, Y be connected regular graphs, with spectra $\{\lambda_i\}, \{\mu_k\}$.*

- (1) *If $\{\lambda_i/\lambda_j\} \cap \{\mu_k/\mu_l\} = \{1\}$ then $G^+(X \circ Y) = G^+(X) \times G^+(Y)$.*
- (2) *If $\{\lambda_i - \lambda_j\} \cap \{\mu_k - \mu_l\} = \{0\}$ then $G^+(X \times Y) = G^+(X) \times G^+(Y)$.*

With these results in hand, it is tempting to try to compute $G^+(X)$ for all vertex-transitive graphs of small order. Here is the result found in [8], up to order 9:

Theorem 3.8. *The vertex-transitive graphs of order ≤ 9 , modulo complementation, and their quantum symmetry groups, are as follows:*

- (1) *Unions of simplices rK_n , having $G^+ = S_n^+ \wr S_r^+$.*
- (2) *Cycles C_n , $n \neq 4$, and the cycles with chords C_8^2, C_9^3 , having $G^+ = D_n$.*
- (3) *The cube, having $G^+ = \mathbb{Z}_2 \times S_4^+$, and $K_3 \times K_3$, having $G^+ = S_3 \wr \mathbb{Z}_2$.*
- (4) *The two squares, $2C_4$, having $G^+ = (\mathbb{Z}_2 \wr \mathbb{Z}_2) \wr \mathbb{Z}_2$.*

We actually have a longer result in [8], going up to order 11, but not covering the Petersen graph. This well-known graph, having 10 vertices, consists of a pentagon with a pentagram inside, with 5 spokes. Our main question, subsequent to [8], is of course:

Problem 3.9. *What is the quantum symmetry group of the Petersen graph?*

Regarding now the proof of Theorem 3.8, this basically follows from Theorem 3.6 and Theorem 3.7, and from a number of “no quantum symmetry” results, coming from Theorem 3.6, from Theorem 3.7, or (e.g. for $K_3 \times K_3$) from a long algebraic computation based on Proposition 3.2. The subject is quite mysterious, for instance because:

$$G(C_n) = G^+(C_n) \iff n \neq 4$$

Indeed, the \implies assertion follows from the discussion after Proposition 3.4. As for the converse, this follows from Proposition 3.2, after a brief computation. See [6].

There is an obvious link here with the following conjecture of Goswami [56]:

Conjecture 3.10. *A compact connected Riemannian manifold cannot have genuine quantum isometries.*

More precisely, Goswami associated in [55] a quantum isometry group $G^+(M)$ to any noncommutative compact Riemannian manifold M in the sense of Connes [45]. In the 0-dimensional case the idea is basically the one in Definition 3.3 above, and in the general case the idea is similar, but more technical, with d being the Laplacian of M .

Goswami was mostly interested in genuine noncommutative manifolds, like the Standard Model one, whose quantum isometries have several potential applications [33]. However, his construction makes sense for any manifold, and leads to the above conjecture.

Back to graphs now, a very interesting example is the n -cube. The symmetry theory of this graph, heavily investigated in [11], can be summarized as follows:

Theorem 3.11. *Let $X = [-1, 1]^n$ be the cube in \mathbb{R}^n , regarded as graph, and let Y be the intersection of X with the n coordinate axes, regarded as a graph formed by n segments.*

- (1) $G(X) = G(Y) = \mathbb{Z}_2 \wr S_n$.
- (2) $G^+(X) = O_n^{-1}$.
- (3) $G^+(Y) = \mathbb{Z}_2 \wr_* S_n^+$

In this statement the group in (1) is the hyperoctahedral group H_n . The quantum group in (2) is a Drinfeld-Jimbo twist at $q = -1$. This twist has the same fusion rules as O_n , so the operation $H_n \rightarrow O_n^{-1}$ is not a “liberation”, in the sense of Theorem 2.10.

As for the quantum group in (3), which obviously looks as the good generalization of H_n , this is called the hyperoctahedral quantum group, and is denoted H_n^+ .

This fact that $H_n \rightarrow H_n^+$ is indeed a “liberation” in the probabilistic sense comes from the following result, once again from [11]:

Theorem 3.12. *The asymptotic laws of truncated characters $\chi_t = \sum_{i=1}^{[tn]} u_{ii}$ are:*

- (1) *For H_n : the Bessel law, $b_t = \text{law}(\alpha - \beta)$, where $\alpha, \beta \sim p_{t/2}$ are independent.*
- (2) *For H_n^+ : the free Bessel law, $\beta_t = \text{law}(\alpha - \beta)$, where $\alpha, \beta \sim \pi_{t/2}$ are free.*

Observe the similarity with Theorem 2.10. The terminology comes from the fact that the density of b_t is given by a Bessel function of the first kind:

$$b_t = e^{-t} \sum_{k=-\infty}^{\infty} \left(\sum_{p=0}^{\infty} \frac{(t/2)^{|k|+2p}}{(|k|+p)!p!} \right) \delta_k$$

As for the measure β_t , this has remarkable properties as well. For instance the odd moments vanish, and the even moments involve the Fuss-Narayana numbers:

$$\int x^{2k} d\beta_t(x) = \sum_{b=1}^k \frac{1}{b} \binom{k-1}{b-1} \binom{2k}{b-1} t^b$$

At the level of diagrams, the partitions as in Theorem 2.3 are those having even blocks, and the algebra as in Theorem 2.7 is the Fuss-Catalan algebra of Bisch and Jones [39].

These results were subsequently explored and generalized to the case of the complex reflection groups $H_n^s = \mathbb{Z}_s \wr S_n$ and their free versions $H_n^{s+} = \mathbb{Z}_s \wr_* S_n^+$. See [7], [30].

A general diagrammatic formalism, covering the quantum groups S_n, S_n^+, H_n, H_n^+ , as well as many other examples, such as the orthogonal group O_n and its free analogue O_n^+ constructed by Wang in [83], was developed in [29], then in [23], [24], [25]:

Definition 3.13. *A compact quantum group $S_n \subset G \subset O_n^+$ is called “easy” if*

$$\text{Hom}(u^{\otimes k}, u^{\otimes l}) = \text{span}(T_\pi | \pi \in D(k, l))$$

for certain sets of partitions $D(k, l) \subset P(k, l)$.

This special class of quantum groups, very useful for unifying certain combinatorial or probabilistic results, was further studied in [46], [47], and in [73].

A promising direction here comes from noncommutative geometry. First, the quantum isometry groups of 0-dimensional manifolds, generalizing the graph construction, were investigated in [34]. In the higher dimensional case, a lot of interesting examples come from discrete group duals [35]. The recent results in [28] show that some of these latter quantum isometry groups are “super-easy”, in the sense that they appear as above, but with the representation $\pi \rightarrow T_\pi$ altered by an involution $J : \mathbb{C}^n \rightarrow \mathbb{C}^n$.

Back now to S_n^+ , the free hypergeometric law of parameters (n, m, N) is the law of:

$$X(n, m, N) = \sum_{i=1}^n \sum_{j=1}^m u_{ij} \in C(S_N^+)$$

The terminology comes from the fact that the variable $X'(n, m, N)$, defined as above, but over the algebra $C(S_N)$, follows a hypergeometric law of parameters (n, m, N) .

Theorem 3.14. *The moments of $X(n, n, n^2)$ are given by*

$$\int X(n, n, n^2)^k dx = \frac{n^k}{(n+1)^k} \cdot \frac{q+1}{q-1} \cdot \frac{1}{k+1} \sum_{r=-k-1}^{k+1} (-1)^r \binom{2k+2}{k+r+1} \frac{r}{1+q^r}$$

where $q \in [-1, 0)$ is given by $q + q^{-1} = -n$.

The proof of this result, from [14], and heavily based on [21], first relies on a purely algebraic twisting result, followed by a monoidal equivalence:

$$(S_{n^2}^+)^r = PO_n^+ \sim SO_3^q$$

The remaining computation can be done by using a number of delicate manipulations involving Askey-Wilson polynomials [3], and leads to the above formula.

These computations are of great interest, for instance in connection with the Atiyah conjecture for quantum groups [43], and with the free spheres introduced in [26].

We would like to end this section with a key algebraic question:

Problem 3.15. *Is the inclusion $S_n \subset S_n^+$ maximal?*

The answer here is yes at $n = 1, 2, 3$, and also yes at $n = 4$, due to the results in [9]. It is also yes in the “easy” case, as shown in [23]. There are some similarities here with a recent result in [13], stating that a certain related inclusion $O_n \subset O_n^*$ is maximal.

4. HADAMARD MATRICES

A complex Hadamard matrix is a matrix $H \in M_n(\mathbb{C})$ whose entries are on the unit circle, $|H_{ij}| = 1$, and whose rows are pairwise orthogonal.

It follows from definitions that the columns of H are pairwise orthogonal as well.

The link with the quantum permutation groups is immediate. Let H_1, \dots, H_n be the rows of H , regarded as invertible elements of the algebra \mathbb{C}^n .

Proposition 4.1. *For an Hadamard matrix $H \in M_n(\mathbb{C})$, the rank one projections*

$$P_{ij} = \text{Proj}(H_i/H_j)$$

form a magic unitary matrix $P \in M_n(A)$, where $A = M_n(\mathbb{C})$.

Indeed, the orthogonality condition on the rows of P follows from the fact that the corresponding vectors form an orthogonal basis of \mathbb{C}^n :

$$\left\langle \frac{H_i}{H_j}, \frac{H_i}{H_k} \right\rangle = \sum_r \bar{H}_{jr} H_{kr} = \delta_{jk}$$

As for the orthogonality condition on the columns, this follows from a similar computation. According now to Definition 1.4 above, we have:

Proposition 4.2. *Associated to any Hadamard matrix $H \in M_n(\mathbb{C})$ is the morphism*

$$\pi_H : C(S_n^+) \rightarrow M_n(\mathbb{C})$$

given by $u_{ij} \rightarrow \text{Proj}(H_i/H_j)$, where H_1, \dots, H_n are the rows of H .

We will be back to this morphism in section 5 below. The idea there will be that associated to π_H is a quantum group $G \subset S_n^+$, describing the “symmetries” of H .

For the moment, let us try to understand what the examples are. First, we have the Fourier matrix, based on the root of unity $w = e^{2\pi i/n}$:

$$F_n = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & w & w^2 & \dots & w^{n-1} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & w^{n-1} & w^{2(n-1)} & \dots & w^{(n-1)^2} \end{pmatrix}$$

This matrix is the matrix of the Fourier transform over \mathbb{Z}_n . There are many other examples of Hadamard matrices, usually coming from finite groups, or other discrete structures. See [76]. An interesting observation, due to Popa, is that a unitary $U \in U_n$ satisfies $\Delta \perp U\Delta U^*$ if and only if $H = \sqrt{n} \cdot U$ is Hadamard. See [72].

In order to present now some classification results, we use the following definitions:

Definition 4.3. *Let H, K be two complex Hadamard matrices.*

- (1) H is called *dephased* if its first row and column consist only of 1's.
- (2) H, K are called *equivalent* if one can pass from one to the other by permuting the rows or columns, or by multiplying them by complex numbers of modulus 1.

Observe that any complex Hadamard matrix can be supposed to be in dephased form, up to the above equivalence relation. With a few exceptions, we will always do so.

At $n = 2, 3$ the Fourier matrix is the only one, up to equivalence. At $n = 4$ we have the following family, depending on a parameter on the unit circle, $q \in \mathbb{T}$:

$$F_4^q = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & q & -q \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -q & q \end{pmatrix}$$

Observe that at $q = 1$ we have $F_4^q = F_2 \otimes F_2$. Observe also that at $q = -1$ we have $F_4^q \simeq F_2 \otimes F_2$, and that at $q = \pm i$ we have $F_4^q \simeq F_4$. We will come back a bit later to these observations, with a discussion regarding the deformations of F_n .

The following remarkable result is due to Haagerup [57].

Theorem 4.4. *The only Hadamard matrices at $n \leq 5$ are F_2, F_3, F_4^q, F_5 .*

At $n = 6$ we have several deformations of $F_6 = F_2 \otimes F_3 = F_3 \otimes F_2$. These matrices appear as particular cases of the following general construction, due to Diță [51]:

Definition 4.5. *The Diță deformation of a tensor product $H \otimes K \in M_{nm}(\mathbb{C})$, with matrix of parameters $L \in M_{m \times n}(\mathbb{T})$, is $H \otimes_L K = (H_{ij} L_{aj} K_{ab})_{ia, jb}$.*

As a first observation, by using the above equivalence relation, we can always assume that L is “dephased”, in the sense that its first row and column consist of 1's.

The basic example comes from deforming the tensor product $F_2 \otimes F_2$, with the matrix of parameters $L = \begin{pmatrix} 1 & 1 \\ 1 & q \end{pmatrix}$. We obtain in this way the above matrix F_4^q :

$$F_4^q = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 1 \\ 1 & q \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

As already mentioned, this matrix can be thought of as being a “deformation” of F_4 , because at $q = \pm i$ we have $F_4^q \simeq F_4$. More generally, we have:

Proposition 4.6. $F_{nm} = F_n \otimes_L F_m$, where $L = (w^{(a-1)(j-1)})_{aj}$ with $w = e^{2\pi i/nm}$.

Let us go back now to the enumeration of Hadamard matrices of small order. At $n = 6$ we have the (dephased) Diță deformations of $F_2 \otimes F_3$ and $F_3 \otimes F_2$, given by:

$$F_6^{(rs)} = F_2 \otimes \begin{pmatrix} 1 & 1 \\ 1 & r \\ 1 & s \end{pmatrix} F_3, \quad F_6^{(rs)} = F_3 \otimes \begin{pmatrix} 1 & 1 & 1 \\ 1 & r & s \end{pmatrix} F_2$$

Here r, s are two parameters on the unit circle, $r, s \in \mathbb{T}$. In matrix form:

$$F_6^{(r)} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & j & j^2 & r & jr & j^2r \\ 1 & j^2 & j & s & j^2s & js \\ 1 & 1 & 1 & -1 & -1 & -1 \\ 1 & j & j^2 & -r & -jr & -j^2r \\ 1 & j^2 & j & -s & -j^2s & -js \end{pmatrix}, \quad F_6^{(rs)} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & r & -r & s & -s \\ 1 & 1 & j & j & j^2 & j^2 \\ 1 & -1 & jr & -jr & j^2s & -j^2s \\ 1 & 1 & j^2 & j^2 & j & j \\ 1 & -1 & j^2r & -j^2r & js & -js \end{pmatrix}$$

There are many other examples at $n = 6$, and we will restrict attention to a certain special class of matrices, defined as follows:

Definition 4.7. *A complex Hadamard matrix is called regular if all the scalar products between distinct rows decompose as sums of cycles.*

Here by ‘‘cycle’’ we mean a sum of p -roots of unity, rotated by a scalar $\lambda \in \mathbb{T}$:

$$C = \sum_{k=1}^p \lambda e^{2k\pi i/p}$$

Besides the above matrices, we have the Haagerup and Tao matrices [57], [77]:

$$H^q = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & i & i & -i & -i \\ 1 & i & -1 & -i & q & -q \\ 1 & i & -i & -1 & -q & q \\ 1 & -i & \bar{q} & -\bar{q} & i & -1 \\ 1 & -i & -\bar{q} & \bar{q} & -1 & i \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & j & j & j^2 & j^2 \\ 1 & j & 1 & j^2 & j^2 & j \\ 1 & j & j^2 & 1 & j & j^2 \\ 1 & j^2 & j^2 & j & 1 & j \\ 1 & j^2 & j & j^2 & j & 1 \end{pmatrix}$$

The following result was proved in [16]:

Theorem 4.8. *The only regular matrices at $n = 6$ are $F_6^{(r)}$, $F_6^{(rs)}$, H^q , T .*

In the non-regular case, the general classification problem is open at $n = 6$. See [75].

One interesting example at $n = 6$, with circulant structure, is the Björck-Fröberg matrix [41], built by using one of the two roots of $a^2 - (1 - \sqrt{3})a + 1 = 0$:

$$BF = \begin{pmatrix} 1 & ia & -a & -i & -\bar{a} & i\bar{a} \\ i\bar{a} & 1 & ia & -a & -i & -\bar{a} \\ -\bar{a} & i\bar{a} & 1 & ia & -a & -i \\ -i & -\bar{a} & i\bar{a} & 1 & ia & -a \\ -a & -i & -\bar{a} & i\bar{a} & 1 & ia \\ ia & -a & -i & -\bar{a} & i\bar{a} & 1 \end{pmatrix}$$

In a remarkable paper [31], Beauchamp and Nicoara classified all self-adjoint Hadamard matrices at $n = 6$. Their proof, and the proof of Theorem 4.8 above too, is heavily

combinatorial. At $n = 7$ we have the following matrix, discovered by Petrescu [71]:

$$P^q = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & qw & qw^4 & w^5 & w^3 & w^3 & w \\ 1 & qw^4 & qw & w^3 & w^5 & w^3 & w \\ 1 & w^5 & w^3 & \bar{q}w & \bar{q}w^4 & w & w^3 \\ 1 & w^3 & w^5 & \bar{q}w^4 & \bar{q}w & w & w^3 \\ 1 & w^3 & w^3 & w & w & w^4 & w^5 \\ 1 & w & w & w^3 & w^3 & w^5 & w^4 \end{pmatrix}$$

Here $w = e^{2\pi i/6}$. This matrix, a non-trivial deformation of prime order, was found by using a computer program, and came as a big surprise at the time of [71].

We can see from the above examples that, at least in the regular case, the roots of unity play a key role in the construction of complex Hadamard matrices.

Definition 4.9. *The level of a complex Hadamard matrix $H \in M_n(\mathbb{C})$ is the smallest number $l \in \{2, 3, \dots, \infty\}$ such that all the entries of H are l -roots of unity.*

Here we agree that a root of unity of order $l = \infty$ is simply an element on the unit circle. It is convenient to introduce as well the following notions:

Definition 4.10. *The Butson class $H_n(l)$ consists of Hadamard matrices in $M_n(\mathbb{C})$ having as entries the l -th roots of unity. In particular:*

- (1) $H_n(2)$ is the set of all $n \times n$ real Hadamard matrices.
- (2) $H_n(l)$ is the set of $n \times n$ Hadamard matrices of level $l|l$.
- (3) $H_n(\infty)$ is the set of all $n \times n$ complex Hadamard matrices.

As a first example, let $H \in H_n(2)$, with $n \geq 3$. By using the equivalence relation, we may assume that the first three rows look blockwise as follows:

$$H = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

Now let a, b, c, d be the lengths of the blocks in the third row. The orthogonality relations between the first three rows give $a + b = c + d$, $a + c = b + d$ and $a + d = b + c$, so we have $a = b = c = d$, and we can conclude that we have $4|n$.

The Hadamard conjecture states that the converse is true:

Conjecture 4.11 (Hadamard conjecture). *If $4|n$ then $H_n(2) \neq \emptyset$.*

This question, going back to the 19th century, is reputed to be of remarkable difficulty. The numeric verification so far goes up to $n = 664$. See [63].

This conjecture has as well an analytic interpretation, that we would like now to explain. Pick any $U \in U_n$ and apply the Cauchy-Schwarz inequality to its entries:

$$\sum_{i,j=1}^n 1 \cdot |U_{ij}| \leq \left(\sum_{i,j=1}^n 1^2 \right)^{1/2} \left(\sum_{i,j=1}^n |U_{ij}|^2 \right)^{1/2}$$

We recognize at left the 1-norm of U . The right term being $n\sqrt{n}$, we get:

$$\|U\|_1 \leq n\sqrt{n}$$

The point now is that the equality holds when the numbers $|U_{ij}|$ are proportional, and since the sum of squares of these numbers is n^2 , we conclude that we have equality if and only if $|U_{ij}| = 1/\sqrt{n}$, i.e. if and only if $H = \sqrt{n} \cdot U$ is Hadamard. Thus we have:

Proposition 4.12. *For $U \in U_n$ we have $\|U\|_1 \leq n\sqrt{n}$, with equality if and only if $U = H/\sqrt{n}$, for a certain complex Hadamard matrix H .*

Now since the maximum of any positive function, and in particular of $U \rightarrow \|U\|_1$, can be obtained from moments, this suggests considering the following type of integral:

$$I_G = \left(\int_G \|U\|_1^k dU \right)^{1/k}$$

More precisely, the Hadamard conjecture can be restated as follows:

$$4|n \implies \lim_{k \rightarrow \infty} I_{O_n} = n\sqrt{n}$$

This observation was made in [19], and a number of related results were obtained there. For instance, it was shown that the Hadamard conjecture is equivalent to:

$$4|n \implies \lim_{k \rightarrow \infty} I_{O_n} \geq n\sqrt{n} - \frac{1}{n\sqrt{n}}$$

This estimate is almost optimal, so the Hadamard conjecture asks for the computation of I_{O_n} with very high accuracy. In the general context of O_n integration, where most results are in the $n \rightarrow \infty$ limit, this kind of problematics is quite new. See [20].

We have the following theoretical questions, regarding the above integrals:

Problem 4.13. *Let $I_G = \left(\int_G \|U\|_1^k dU \right)^{1/k}$.*

- (1) *Does the first-order term of I_{O_n} count $|H_n(2)|$?*
- (2) *Is it possible to generalize this to any $l < \infty$?*
- (3) *What does the first-order term of I_{U_n} count?*

These questions are of course very vague, because we don't know what "first-order term" should mean. As an answer to the third question, we would expect of course some very simple numeric invariant of the complex Hadamard matrix manifold $H_n(\infty)$.

Let us go back now to the arbitrary sets of Butson matrices, $H_n(l)$ with $l < \infty$.

Theorem 4.14. *We have the following list of obstructions:*

- (1) *Lam-Leung:* If $H_n(l) \neq \emptyset$ and $l = p_1^{a_1} \dots p_s^{a_s}$ then $n \in p_1\mathbb{N} + \dots + p_s\mathbb{N}$.
- (2) *de Launey:* If $H_n(l) \neq \emptyset$ then there is $d \in \mathbb{Z}[e^{2\pi i/l}]$ such that $|d|^2 = n^n$.
- (3) *Sylvester:* If $H_n(2) \neq \emptyset$ then $n = 2$ or $4|n$.
- (4) *Haagerup:* If $H_5(l) \neq \emptyset$ then $5|l$.

In this statement (1) follows from [66], (2), where $d = \det H$, was studied in [67], (3) was already explained, and (4) comes from Theorem 4.4 above.

The proof of the Sylvester obstruction, presented above, can be adapted for higher values of l , and leads to the following supplementary ‘‘Sylvester obstructions’’:

- (1) If $H_n(l) \neq \emptyset$ and $n = p + 2$ with $p \geq 3$ prime, then $l \neq 2p^b$.
- (2) If $H_n(l) \neq \emptyset$ and $n = 2q$ with $p > q \geq 3$ primes, then $l \neq 2^a p^b$.

These results were proved in [16]. The point now is that, with these results in hand, the case $n \leq 10, l \leq 14$ is fully covered. We have the following table from [16], describing for each n, l either an explicit matrix in $H_n(l)$, or an obstruction which applies:

$n \setminus l$	2	3	4	5	6	7	8	9	10	11	12	13	14
2	F_2	\circ	F_2	\circ	F_2	\circ	F_2	\circ	F_2	\circ	F_2	\circ	F_2
3	\circ	F_3	\circ	\circ	F_3	\circ	\circ	F_3	\circ	\circ	F_3	\circ	\circ
4	F_{22}	\circ	F_{22}	\circ	F_{22}	\circ	F_{22}	\circ	F_{22}	\circ	F_{22}	\circ	F_{22}
5	\circ	\circ	\circ	F_5	\circ_l	\circ	\circ	\circ	F_5	\circ	\circ_h	\circ	\circ
6	\circ_s	T	H^1	\circ	T	\circ	H^1	T	\circ_s	\circ	T	\circ	\circ_s
7	\circ	\circ	\circ	\circ	P^1	F_7	\circ	\circ	\circ_s	\circ	P^1	\circ	F_7
8	F_{222}	\circ	F_{222}	\circ	F_{222}	\circ	F_{222}	\circ	F_{222}	\circ	F_{222}	\circ	F_{222}
9	\circ	F_{33}	\circ	\circ	F_{33}	\circ	\circ	F_{33}	W	\circ	F_{33}	\circ	\circ_s
10	\circ_s	\circ	X	Y	Z	\circ	X	\circ	F_{10}	\circ	X	\circ	\circ

Here $\circ, \circ_l, \circ_s, \circ_h$ denote the various obstructions in Theorem 4.14 and in the discussion afterwards, for $k_1, \dots, k_s \in \{2, 3\}$ we use the notation $F_{k_1 \dots k_s} = F_{k_1} \otimes \dots \otimes F_{k_s}$, and X, Y, Z, W are matrices constructed by a computer, and presented in [16].

Question 4.15. *What happens at $n = 11$? And, what happens at $l = 15$?*

We refer here to [16] for technical details.

Finally, let us mention the following key conjecture, from [16].

Conjecture 4.16. *Any Butson matrix is regular.*

One difficulty in approaching this conjecture lies in the fact that the vanishing sums of roots of unity have a quite complicated combinatorics, cf. [66].

5. THE CORRESPONDENCE

We have seen in the previous section that the complex Hadamard matrices, while strongly related to the quantum permutation groups via the magic condition, seem rather to belong to another planet. There is no obvious relation between the substantial mass of combinatorics presented in sections 1-3 above, and the substantial mass of combinatorics presented in section 4 above. So, the following statement might seem a bit strange:

Fact 5.1. *The symmetries of a complex Hadamard matrix $H \in M_n(\mathbb{C})$ are described by a quantum permutation group $G \subset S_n^+$. Conversely, each quantum permutation group $G \subset S_n^+$ is the symmetry group of an Hadamard-type object $H \in \tilde{M}_n(\mathbb{C})$.*

In this section we will try to explain why this fact is true. The fact that the known combinatorics of quantum permutation groups remains quite far from the known combinatorics of complex Hadamard matrices will be discussed in the next section. We will propose there some conjectural answers to the question, making the above fact a potentially useful tool, in the study of both quantum permutations and Hadamard matrices.

Let us begin with the notion of Hopf image, recently axiomatized in [10]:

Definition 5.2. *The Hopf image of A by a representation $\pi : A \rightarrow M_n(\mathbb{C})$ is the smallest Hopf C^* -algebra A' producing a factorization $\pi : A \rightarrow A' \rightarrow M_n(\mathbb{C})$.*

In order to understand this notion, let $A = C^*(\Gamma)$. Then π must come from a unitary group representation $\tilde{\pi} : \Gamma \rightarrow U_n$, and we have $A' = C^*(\Lambda)$, where $\Lambda = \tilde{\pi}(\Gamma)$.

In this computation Γ was of course a usual discrete group. In the general case, i.e. when Γ is a discrete quantum group, it is only known that the Hopf image exists, and is unique [10]. But, of course, the discrete quantum group point of view is very useful.

Here are a few more definitions from [10], based on the same philosophy:

Definition 5.3. *Let A be a Hopf C^* -algebra.*

- (1) *A representation $\pi : A \rightarrow M_n(\mathbb{C})$ is called inner faithful if $A = A'$.*
- (2) *A is called inner linear if it has an inner faithful representation.*

Observe that with $A = C^*(\Gamma)$, and with the above notations, π is inner faithful if and only if $\tilde{\pi}$ is faithful. Also, A is inner linear if and only if Γ is linear.

These notions, coming from [4], [27], [16], were axiomatized and studied in [10]. A number of key examples were later on studied by Andruskiewitsch and Bichon in [2].

Some useful supplementary tools, of functional analytic nature, are expected to come from the idempotent state theory of Franz and Skalski [54].

We can apply these constructions to the representation π_H in Proposition 4.2:

Definition 5.4. *Associated to any complex Hadamard matrix $H \in M_n(\mathbb{C})$ is the quantum permutation group $G \subset S_n^+$ given by the fact that $C(G)$ is the Hopf image of π_H .*

In other words, if we go back to the definition of both π_H and of the Hopf image, $C(G)$ is the smallest Hopf C^* -algebra producing the following factorization:

$$\begin{aligned} C(S_n^+) &\rightarrow C(G) \rightarrow M_n(\mathbb{C}) \\ u_{ij} &\rightarrow Proj(H_i/H_j) \end{aligned}$$

Equivalently, we can say that this representation π_H should correspond to a unitary quantum group representation $\tilde{\pi}_H : \widehat{S}_n^+ \rightarrow U_n^+$, and G is simply given by:

$$\widehat{G} = \tilde{\pi}_H(\widehat{S}_n^+)$$

This latter description of the correspondence $H \rightarrow G$, while being quite heuristic, starts to justify the first claim in Fact 5.1, namely that “ G encodes the symmetries of H ”.

Proposition 5.5. *The construction $H \rightarrow G$ has the following properties:*

- (1) *For $H = F_n$ we obtain $G = \mathbb{Z}_n$.*
- (2) *For $H = H' \otimes H''$ we obtain $G = G' \times G''$.*

The first assertion follows indeed from the definition of the Hopf image, because the representation $C(S_n^+) \rightarrow M_n(\mathbb{C})$ associated to F_n is circulant, and hence factorizes through $C(\mathbb{Z}_n)$. Now since this factorization cannot be further factorized, we obtain $G = \mathbb{Z}_n$.

As for the second assertion, this follows as well from definitions. See [27], [16].

More generally, we have the following result, once again from [27], [16]:

Theorem 5.6. *For an Hadamard matrix H , the following are equivalent:*

- (1) *G is a classical group.*
- (2) *\widehat{G} is a classical group.*
- (3) *$G = \mathbb{Z}_{n_1} \times \dots \times \mathbb{Z}_{n_k}$, for some numbers n_1, \dots, n_k .*
- (4) *$H = F_{n_1} \otimes \dots \otimes F_{n_k}$, for some numbers n_1, \dots, n_k .*

We recall now from Definition 4.5 above that the Diță deformation of a tensor product $H \otimes K \in M_{nm}(\mathbb{C})$, with matrix of parameters $L \in M_{m \times n}(\mathbb{T})$, is:

$$H \otimes_L K = (H_{ij} L_{aj} K_{ab})_{ia, jb}$$

We have already seen in section 4 that this notion is a key one. For instance Theorem 4.4 tells us that at $n = 4$ the only Hadamard matrices are the Diță deformations of $F_2 \otimes F_2$, and Theorem 4.8 tells us that the only regular matrices at $n = 6$ are the Haagerup and Tao matrices H^q, T , and the Diță deformations of $F_6 = F_2 \otimes F_3 = F_3 \otimes F_2$.

The symmetries of $H \otimes_L K$, even for very simple values of H, K , vary a lot, and in a very subtle way, with the matrix of parameters L . The only general result here is:

Theorem 5.7. *We have $G(H \otimes_L K) \subset G(K) \wr_* G(H)$.*

This result, from [16], marks the end of the algebraic theory. We will come back in section 6 below with a conjectural discussion of the problem.

For the moment, let us just record a difficult question:

Problem 5.8. *What is $G(F_n \otimes_L F_m)$?*

Here are a few more questions, for the most of algebraic nature:

Problem 5.9. *Let H be a complex Hadamard matrix.*

- (1) *When is G finite?*
- (2) *When is \widehat{G} amenable?*
- (3) *Can we have $G = S_n^+$?*

All these questions look equally very difficult. Question (3) is due to Jones [61]. Let us discuss now the representation theory of G . The result here, from [16], is as follows:

Theorem 5.10. *We have $T \in \text{Hom}(u^{\otimes k}, u^{\otimes l})$ if and only if $T^\circ G^{k+2} = G^{l+2} T^\circ$, where:*

- (1) $T^\circ = id \otimes T \otimes id$.
- (2) $G_{ia}^{jb} = \sum_{k=1}^n H_{ik} \bar{H}_{jk} \bar{H}_{ak} H_{bk}$.
- (3) $G_{i_1 \dots i_k, j_1 \dots j_k}^k = G_{i_k i_{k-1}}^{j_k j_{k-1}} \cdots G_{i_2 i_1}^{j_2 j_1}$.

The proof in [16] uses a quite simple Tannakian argument, further developed in [10], stating that when we have an inner faithful representation $u_{ij} \rightarrow P_{ij}$, then:

$$\text{Hom}(u^{\otimes k}, u^{\otimes l}) = \text{Hom}(P^{\otimes k}, P^{\otimes l})$$

Here the spaces on the right are defined exactly as those on the left, namely:

$$\text{Hom}(P^{\otimes k}, P^{\otimes l}) = \{T : (\mathbb{C}^n)^{\otimes k} \rightarrow (\mathbb{C}^n)^{\otimes l} \mid TP^{\otimes k} = P^{\otimes l}T\}$$

This result makes the link with the planar algebra approach of Jones [61]:

Theorem 5.11. *Let H be a complex Hadamard matrix.*

- (1) *The planar algebra associated to H is given by $P_k = \text{Fix}(u^{\otimes k})$.*
- (2) *The quantum invariants of H are the moments of $\chi = \text{Tr}(u)$.*
- (3) *The Poincaré series of H is the Stieltjes transform of $\text{law}(\chi)$.*

Indeed, (1) follows Theorem 5.10 at $k = 0$, because the description there of the space $\text{Fix}(u^{\otimes l})$ coincides with the description in [61] of the space P_l . As for (2,3), these are simply quantum group translations of Jones' notions in [61], namely:

$$c_k = \dim(P_k), \quad f(z) = \sum_{k=0}^{\infty} c_k z^k$$

More precisely, using the results of Woronowicz in [85], we get as claimed:

$$c_k = \int_G \chi^k, \quad f(x) = \int_G \frac{1}{1 - z\chi}$$

Summarizing, the above results fully justify the first claim in Fact 5.1, namely that “ G is the symmetry group of H ”. The story here can be actually made longer, by talking about commuting squares [72], spin models [60] and subfactors [59]. See [4].

We should probably mention that, via the various algebraic results presented in the beginning of this section, Theorem 5.11 allows one to recover most of what is known about the Hadamard matrix subfactors, sometimes with some slight enhancements.

Let us go now to the second claim in Fact 5.1, namely that any quantum permutation group $G \subset S_n^+$ is the symmetry group of an Hadamard-type object $H \in \tilde{M}_n(\mathbb{C})$. The situation here is much more complicated, and we only have a categorical result so far:

Theorem 5.12. *There is a Tannaka-Galois correspondence between quantum permutation groups $G \subset S_n^+$ and subalgebras of the spin planar algebra $P \subset Q_n$. In addition:*

- (1) *The quantum symmetry groups of generalized colored graphs correspond in this way to the subalgebras of Q_n generated by a 2-box.*
- (2) *The quantum groups S_n^+, H_n^+ correspond in this way to the canonical copies of the Temperley-Lieb and Fuss-Catalan algebras inside Q_n .*
- (3) *The quantum group associated to a complex Hadamard matrix $H \in M_n(\mathbb{C})$ corresponds in this way to the planar algebra associated to H .*

The Tannaka-Galois correspondence was established in [6], the idea being that the embedding $P \subset Q_n$ could be regarded as a “piece” of the usual Tannakian data, namely “tensor category + Hilbert functor”. By Frobenius duality we can recover the rest of the Tannakian data, and by using Woronowicz’s duality in [86] we obtain the result.

The assertion (1) was equally established in [6], the idea here being simply that the condition $d \in \text{End}(u)$ reads $d \in P_2$, and hence $P = \langle d \rangle$. This result shows that our classification program for graphs, presented in section 3, can be regarded as part of the Bisch-Jones classification program for planar algebras generated by a 2-box [40].

The assertion (2) for S_n^+ follows simply from functoriality: since S_n^+ is the biggest quantum permutation group, it should correspond to the smallest planar subalgebra of Q_n , which is of course the Temperley-Lieb algebra TL_n . Observe that we have here yet another proof of Theorem 2.7, but this time, without any single computation!

As for the assertion (2) for H_n^+ , this follows from Theorem 3.12 above, or rather from the diagrammatic comments following it. See [11]. The interest in this statement comes of course in conjunction with Theorem 3.12, and with the various probabilistic formulae following it: all these formulae concern of course the Fuss-Catalan algebra of Bisch and Jones [39], which was previously not known to have all these combinatorial aspects.

Finally, the assertion (3) follows from Theorem 5.11 above.

Let us go back now to the second claim in Fact 5.1. Given a quantum permutation group $G \subset S_n^+$, we would like to construct an object $H \in \tilde{M}_n(\mathbb{C})$ whose symmetry group is G . This is of course a quite difficult task, and the above categorical result, or the purely algebraic considerations in [16], are just a first step towards an answer.

We believe that the following question can push things here one step further:

Problem 5.13. *What is the noncommutative geometry of a complex Hadamard matrix?*

In other words, associated to any complex Hadamard matrix $H \in M_n(\mathbb{C})$ we would like to have a noncommutative manifold M_H in the sense of Connes [45], modelled on the noncommutative set $M_n = \text{Spec}(M_n(\mathbb{C}))$, with the requirement $G(H) = G^+(M)$.

6. MATRIX MODELS

We have seen in the previous section that associated to any complex Hadamard matrix $H \in M_n(\mathbb{C})$ is a quantum permutation group $G \subset S_n^+$, describing its “symmetries”. The correspondence $H \rightarrow G$ appears to be fully satisfactory at the level of simple examples, and is also compatible with the planar algebra approach of Jones [61].

The challenging problem is to relate the combinatorics of quantum permutation groups, explained in sections 1-3 above, to the combinatorics of complex Hadamard matrices, explained in section 4 above. Our answer here, largely conjectural, is as follows:

Conjecture 6.1. *Consider the correspondence $H \rightarrow G$.*

- (1) *The known combinatorics of G and the known combinatorics of H are not different: they are just different aspects of a single combinatorics.*
- (2) *This single combinatorics appears when looking at the affine families $\{H^q | q \in \mathbb{T}^r\}$, and at the corresponding quantum groups $\{G^q | q \in \mathbb{T}^r\}$.*
- (3) *The unifying theorem is a statement regarding matrix models for G^q, H^q , with the parameter space being a certain subgroup of \mathbb{T}^r , associated to q .*

In other words, our claim is that the quantum permutation groups $G^q = G(H^q)$ with $q \in \mathbb{T}^r$ should behave a bit like the usual Drinfeld-Jimbo deformations [52], [58] in the parameter range $|q| = 1$, with a rich combinatorics depending on the arithmetics of q . And this combinatorics should correspond to the various combinatorial aspects of H^q , i.e. to the Butson obstructions, and to the subtle behavior of the Poincaré series.

Our second claim is that a useful and effective tool for dealing with these difficult questions in the notion of matrix model. The idea here, to be explained in detail later on, is that under suitable assumptions we should have an explicit matrix model of type $C(G^q) \subset C(K_q) \otimes M_n(\mathbb{C})$, where $K_q \subset \mathbb{T}^r$ is a certain abelian group associated to q .

Let us first explain the notion of affine family:

Definition 6.2. *An affine family of complex Hadamard matrices is a family $H^q \in M_n(\mathbb{C})$, depending analytically on a parameter $q \in \mathbb{T}^r$. The family is called:*

- (1) *Regular, if all matrices H^q are regular.*
- (2) *Normalized, if $l(H^1) \leq l(H^q)$, for any $q \in \mathbb{T}^r$.*
- (3) *Arithmetic, if $l(H^q) \leq K \cdot \text{ord}(q)$, for some $K > 0$.*

We use here the notions of regularity and level, introduced in section 4. As an example, the Diţă deformations of Fourier matrices, as well as the Haagerup and Petrescu matrices H^q, P^q , form affine families which are regular, arithmetic, and often normalized.

With a bit of complex analysis, one can prove that any arithmetic regular family is obtained in a similar way, by starting with a Butson matrix, and adding r parameters.

We recall from [69], [76] that the “defect” of a matrix $H \in M_n(\mathbb{C})$ counts the dimension of the component of the Hadamard matrix manifold $H_n(\infty)$ passing through H .

Here is a first question that we have, regarding the affine families:

Problem 6.3. *Is there a notion of “regular defect” of $H \in M_n(\mathbb{C})$, counting the dimension of the maximal regular affine family passing through H ?*

In other words, we are asking here for an explicit formula for the regular defect, in the spirit of those found in [76]. This question is probably quite difficult, due to its relation with Conjecture 4.16 above, stating that any Butson matrix is regular.

Here is another question, probably more elementary:

Problem 6.4. *Is it true that the maximal affine family passing through $F = F_{n_1} \otimes \dots \otimes F_{n_k}$ can be obtained via a sequence of Diţă deformations of F ?*

Normally this should follow from the various defect formulae in [76] and from the fact that the Fourier matrix $F_{n_1 \dots n_k}$ is in the orbit of F , cf. Proposition 4.6.

Let us discuss now the notion of matrix model:

Definition 6.5. *A matrix model for a Hopf C^* -algebra A is a morphism of C^* -algebras $\pi : A \rightarrow L^\infty(X) \otimes B(H)$, where X is a probability space, and H is a finite dimensional Hilbert space, such that $\int a = (\mathbb{E} \otimes \text{tr})\pi(a)$, for any $a \in A$.*

At the level of examples, let us first look at the commutative case. Here the algebras in question are those of the form $A = C(G)$, where G is a usual compact group, and the identity map $\text{id} : C(G) \rightarrow L^\infty(G) \otimes B(\mathbb{C})$ is of course a matrix model.

Let us record now a few general properties of the matrix models.

Proposition 6.6. *Let $\pi : A \rightarrow L^\infty(X) \otimes B(H)$ be a matrix model.*

- (1) *A is amenable in the discrete quantum group sense.*
- (2) *π is injective, i.e. $\pi(a) = 0$ implies $a = 0$.*

Indeed, let $\varphi : A \rightarrow A_{red}$ the projection onto the reduced version, obtained by dividing by the null ideal of the Haar functional. For any $a \in A$ we have:

$$\pi(a) = 0 \implies (\mathbb{E} \otimes \text{tr})\pi(aa^*) = 0 \implies \int aa^* = 0 \implies \varphi(a) = 0$$

Thus we have $\ker \pi \subset \ker \varphi$, so A_{red} appears as quotient of the image algebra $\pi(A) \subset L^\infty(X) \otimes B(H)$. With this observation in hand, we can prove now our assertions:

Indeed, (1) follows from the fact that $L^\infty(X) \otimes B(H)$ is of type I: by the general theory, so must be its subalgebra $\pi(A)$, and also its subquotient A_{red} . But now since A_{red} is of type I, A follows to be amenable, i.e. we have $A = A_{red}$, and we are done.

As for (2), since A is amenable we have $\ker \pi \subset \ker \varphi = \mathbb{C}$, and we are done.

Definition 6.7. *A linear model for a quantum permutation algebra $C(S_n^+) \rightarrow A$ is a continuous matrix model $\pi : A \rightarrow C(K) \otimes M_n(\mathbb{C})$, where $K \subset O_n$ is a compact group, such that the magic entries $U_{ij}^x \in M_n(\mathbb{C})$ are linear functions of $x \in K \subset M_n(\mathbb{C})$.*

In other words, we are making some key assumptions here, both taking into account the number n of points on which A coacts: we assume that our probability space is a compact group $K \subset O_n$, that our Hilbert space is $H = \mathbb{C}^n$, and that the model itself comes from a “linear construction” applied to the embedding $K \subset B(H)$.

The basic example is the model for $C(S_4^+)$ found in [18]. Recall first that the elements of SU_2 are of the following form, with $(x, y, z, t) \in S^3$:

$$m = \begin{pmatrix} x + iy & z + it \\ -z + it & x - iy \end{pmatrix}$$

Observe that $m = xc_1 + yc_2 + zc_3 + tc_4$, where c_i are the Pauli matrices:

$$c_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad c_2 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad c_3 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad c_4 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$$

Now SU_2 acts by conjugation, $m'(p) = mpm^*$, on $\text{span}(c_1, c_2, c_3, c_4) \simeq \mathbb{R}^4$, and we have $m' = \text{diag}(1, m'')$, with $m'' \in SO_3$. Thus we have explicit morphisms $SU_2 \rightarrow SO_3 \subset O_4$.

Theorem 6.8. *We have a linear model $C(S_4^+) \subset C(SO_3) \otimes M_4(\mathbb{C})$ given by $U_{ij}^x = \text{Proj}(c_i x c_j)$, where c_1, c_2, c_3, c_4 are the Pauli matrices.*

This result was found in [18]. The first observation is that the formula of U_{ij}^x in the statement defines a morphism of algebras $C(S_4^+) \rightarrow C(SU_2) \otimes M_4(\mathbb{C})$. Now since U_{ij}^x is invariant under $x \rightarrow -x$, this shows that the target of our morphism is contained in $C(SO_3) \otimes M_4(\mathbb{C})$. Thus we have a representation as in the statement.

As pointed out in [9], this representation is indeed a linear model in the sense of Definition 6.7, when using the above-mentioned embedding $SO_3 \subset O_4$.

The proof in [18] of the equality $\int a = (\mathbb{E} \otimes \text{tr})\pi(a)$ uses the Weingarten formula for the evaluation of the left term, and an explicit computation for the right term.

We have now all ingredients for stating our main conjecture:

Conjecture 6.9. *Let $\{H^q | q \in \mathbb{T}^r\}$ be an affine family of complex Hadamard matrices, and consider the corresponding family of quantum permutation groups $\{G^q | q \in \mathbb{T}^r\}$. Then, under suitable assumptions, there exist compact abelian groups $K_q \subset \mathbb{T}^r$ such that:*

- (1) *We have twisting results of type $G_q = K_q^{-1}$.*
- (2) *We have linear models of type $C(G_q) \subset C(K_q) \otimes M_n(\mathbb{C})$.*

Observe that at $n = 4$ we know from Theorem 5.7 above that we have an inclusion $G(F_4^q) \subset \mathbb{Z}_2 \lambda_* \mathbb{Z}_2 = O_2^{-1}$, so the remaining problem is to find the correct group K_q , by using Theorem 1.12 and Theorem 6.8. This will be explained in a forthcoming paper.

CONCLUSION

We have seen in this paper that the combinatorics of quantum permutation groups $G \subset S_n^+$ shares many common aspects with the combinatorics of complex Hadamard matrices $H \in M_n(\mathbb{C})$. The quite substantial mass of work so far, on quantum groups and Hadamard matrices, goes however in slightly different directions. We can only hope that a further study of the correspondence $H \rightarrow G$ would be fruitful on both sides.

One problem comes from the fact that a lot of work on G has gone into quite delicate probabilistic aspects, whose matrix counterpart is for the moment unknown.

The other problem is that a lot of work on H has gone into the root of unity case, and an abstract notion of quantum permutation groups at roots of unity is still lacking.

We believe that both problems can be solved, and the correspondence $H \rightarrow G$ can be substantially upgraded, in the general context of the matrix model problematics. This idea, which is largely conjectural for the moment, is explained in section 6 above.

Finally, we hope that the present text was readable. The prerequisites for a second reading would be a certain familiarity with Hopf algebras and operator algebras [1], [70], with compact quantum groups [85], [86], and with free probability theory [68], [81]. The reader might also consult the previous survey article on the subject [12].

REFERENCES

- [1] E. Abe, Hopf algebras, Cambridge Univ. Press (1977)
- [2] N. Andruskiewitsch and J. Bichon, Examples of inner linear Hopf algebras, *Rev. Un. Mat. Argentina* **51** (2010), 7–18.
- [3] R.A. Askey and J. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Mem. Amer. Math. Soc.* **54** (1985).
- [4] T. Banica, Hopf algebras and subfactors associated to vertex models, *J. Funct. Anal.* **159** (1998), 243–266.
- [5] T. Banica, Representations of compact quantum groups and subfactors, *J. Reine Angew. Math.* **509** (1999), 167–198.
- [6] T. Banica, Quantum automorphism groups of homogeneous graphs, *J. Funct. Anal.* **224** (2005), 243–280.
- [7] T. Banica, S.T. Belinschi, M. Capitaine and B. Collins, Free Bessel laws, *Canad. J. Math.* **63** (2011), 3–37.
- [8] T. Banica and J. Bichon, Quantum automorphism groups of vertex-transitive graphs of order ≤ 11 , *J. Algebraic Combin.* **26** (2007), 83–105.
- [9] T. Banica and J. Bichon, Quantum groups acting on 4 points, *J. Reine Angew. Math.* **626** (2009), 74–114.
- [10] T. Banica and J. Bichon, Hopf images and inner faithful representations, *Glasg. Math. J.* **52** (2010), 677–703.
- [11] T. Banica, J. Bichon and B. Collins, The hyperoctahedral quantum group, *J. Ramanujan Math. Soc.* **22** (2007), 345–384.
- [12] T. Banica, J. Bichon and B. Collins, Quantum permutation groups: a survey, *Banach Center Publ.* **78** (2007), 13–34.

- [13] T. Banica, J. Bichon, B. Collins and S. Curran, A maximality result for orthogonal quantum groups, *Comm. Algebra*, to appear.
- [14] T. Banica, J. Bichon and S. Curran, Quantum automorphisms of twisted group algebras and free hypergeometric laws, *Proc. Amer. Math. Soc.* **139** (2011), 3961–3971.
- [15] T. Banica, J. Bichon and S. Natale, Finite quantum groups and quantum permutation groups, *Adv. Math.* **229** (2012), 3320–3338.
- [16] T. Banica, J. Bichon and J.-M. Schlenker, Representations of quantum permutation algebras, *J. Funct. Anal.* **257** (2009), 2864–2910.
- [17] T. Banica and B. Collins, Integration over quantum permutation groups, *J. Funct. Anal.* **242** (2007), 641–657.
- [18] T. Banica and B. Collins, Integration over the Pauli quantum group, *J. Geom. Phys.* **58** (2008), 942–961.
- [19] T. Banica, B. Collins and J.-M. Schlenker, On orthogonal matrices maximizing the 1-norm, *Indiana Univ. Math. J.* **59** (2010), 839–856.
- [20] T. Banica, B. Collins and J.-M. Schlenker, On polynomial integrals over the orthogonal group, *J. Combin. Theory Ser. A* **118** (2011), 778–795.
- [21] T. Banica, B. Collins and P. Zinn-Justin, Spectral analysis of the free orthogonal matrix, *Int. Math. Res. Not.* **17** (2009), 3286–3309.
- [22] T. Banica and S. Curran, Decomposition results for Gram matrix determinants, *J. Math. Phys.* **51** (2010), 1–14.
- [23] T. Banica, S. Curran and R. Speicher, Classification results for easy quantum groups, *Pacific J. Math.* **247** (2010), 1–26.
- [24] T. Banica, S. Curran and R. Speicher, Stochastic aspects of easy quantum groups, *Probab. Theory Related Fields* **149** (2011), 435–462.
- [25] T. Banica, S. Curran and R. Speicher, De Finetti theorems for easy quantum groups, *Ann. Probab.* **40** (2012), 401–435.
- [26] T. Banica and D. Goswami, Quantum isometries and noncommutative spheres, *Comm. Math. Phys.* **298** (2010), 343–356.
- [27] T. Banica and R. Nicoara, Quantum groups and Hadamard matrices, *Panamer. Math. J.* **17** (2007), 1–24.
- [28] T. Banica and A. Skalski, Two-parameter families of quantum symmetry groups, *J. Funct. Anal.* **260** (2011), 3252–3282.
- [29] T. Banica and R. Speicher, Liberation of orthogonal Lie groups, *Adv. Math.* **222** (2009), 1461–1501.
- [30] T. Banica and R. Vergnioux, Fusion rules for quantum reflection groups, *J. Noncommut. Geom.* **3** (2009), 327–359.
- [31] K. Beauchamp and R. Nicoara, Orthogonal maximal abelian $*$ -subalgebras of the 6×6 matrices, *Linear Algebra Appl.* **428** (2008), 1833–1853.
- [32] H. Bercovici and V. Pata, Stable laws and domains of attraction in free probability theory, *Ann. of Math.* **149** (1999), 1023–1060.
- [33] J. Bhowmick, F. D’Andrea and L. Dabrowski, Quantum isometries of the finite noncommutative geometry of the standard model, *Comm. Math. Phys.* **307** (2011), 101–131.
- [34] J. Bhowmick, D. Goswami and A. Skalski, Quantum isometry groups of 0-dimensional manifolds, *Trans. Amer. Math. Soc.* **363** (2011), 901–921.
- [35] J. Bhowmick and A. Skalski, Quantum isometry groups of noncommutative manifolds associated to group C^* -algebras, *J. Geom. Phys.* **60** (2010), 1474–1489.
- [36] J. Bichon, Free wreath product by the quantum permutation group, *Alg. Rep. Theory* **7** (2004), 343–362.

- [37] J. Bichon, Algebraic quantum permutation groups, *Asian-Eur. J. Math.* **1** (2008), 1–13.
- [38] J. Bichon, A. De Rijdt and S. Vaes, Ergodic coactions with large multiplicity and monoidal equivalence of quantum groups, *Comm. Math. Phys.* **262** (2006), 703–728.
- [39] D. Bisch and V.F.R. Jones, Algebras associated to intermediate subfactors, *Invent. Math.* **128** (1997), 89–157.
- [40] D. Bisch and V.F.R. Jones, Singly generated planar algebras of small dimension, *Duke Math. J.* **101** (2000), 41–75.
- [41] G. Björck and R. Fröberg, A faster way to count the solutions of inhomogeneous systems of algebraic equations, with applications to cyclic n -roots, *J. Symbolic Comput.* **12** (1991), 329–336.
- [42] M. Brannan, Approximation properties for free orthogonal and free unitary quantum groups, *J. Reine Angew. Math.*, to appear.
- [43] B. Collins, J. Härtel and A. Thom, Homology of free quantum groups, *C. R. Math. Acad. Sci. Paris* **347** (2009), 271–276.
- [44] B. Collins and P. Śniady, Integration with respect to the Haar measure on unitary, orthogonal and symplectic groups, *Comm. Math. Phys.* **264** (2006), 773–795.
- [45] A. Connes, Noncommutative geometry, Academic Press (1994).
- [46] S. Curran, A characterization of freeness by invariance under quantum spreading, *J. Reine Angew. Math.*, to appear.
- [47] S. Curran and R. Speicher, Quantum invariant families of matrices in free probability, *J. Funct. Anal.* **261** (2011), 897–933.
- [48] K. De Commer, On projective representations for compact quantum groups, *J. Funct. Anal.* **260** (2011), 3596–3644.
- [49] P. Di Francesco, O. Golinelli and E. Guitter, Meanders and the Temperley-Lieb algebra, *Comm. Math. Phys.* **186** (1997), 1–59.
- [50] P. Diaconis and M. Shahshahani, On the eigenvalues of random matrices, *J. Applied Probab.* **31** (1994), 49–62.
- [51] P. Diță, Some results on the parametrization of complex Hadamard matrices, *J. Phys. A* **37** (2004), 5355–5374.
- [52] V. Drinfeld, Quantum groups, Proc. ICM Berkeley (1986), 798–820.
- [53] P. Fima, Property T for discrete quantum groups, *Internat. J. Math.* **21** (2010), 47–65.
- [54] U. Franz and A. Skalski, On idempotent states on quantum groups, *J. Algebra* **322** (2009), 1774–1802.
- [55] D. Goswami, Quantum group of isometries in classical and noncommutative geometry, *Comm. Math. Phys.* **285** (2009), 141–160.
- [56] D. Goswami, Rigidity of action of compact quantum groups, [arxiv:1106.5107](https://arxiv.org/abs/1106.5107).
- [57] U. Haagerup, Orthogonal maximal abelian $*$ -subalgebras of the $n \times n$ matrices, in “Operator algebras and quantum field theory”, International Press (1997), 296–323.
- [58] M. Jimbo, A q -difference analog of $U(g)$ and the Yang-Baxter equation, *Lett. Math. Phys.* **10** (1985), 63–69.
- [59] V.F.R. Jones, Index for subfactors, *Invent. Math.* **72** (1983), 1–25.
- [60] V.F.R. Jones, On knot invariants related to some statistical mechanical models, *Pacific J. Math.* **137** (1989), 311–334.
- [61] V.F.R. Jones, Planar algebras I, [arxiv:math/9909027](https://arxiv.org/abs/math/9909027).
- [62] D. Kazhdan and H. Wenzl, Reconstructing monoidal categories, *Adv. in Soviet Math.* **16** (1993), 111–136.
- [63] H. Kharaghani and B. Tayfeh-Rezaie, A Hadamard matrix of order 428, *J. Combin. Des.* **13** (2005), 435–440.

- [64] C. Köstler, R. Speicher, A noncommutative de Finetti theorem: invariance under quantum permutations is equivalent to freeness with amalgamation, *Comm. Math. Phys.* **291** (2009), 473–490.
- [65] D. Kyed, L^2 -homology for compact quantum groups, *Math. Scand.* **103** (2008), 111–129.
- [66] T.Y. Lam and K.H. Leung, On vanishing sums of roots of unity, *J. Algebra* **224** (2000), 91–109.
- [67] W. de Launey, On the non-existence of generalized weighing matrices, *Ars Combin.* **17** (1984), 117–132.
- [68] A. Nica and R. Speicher, Lectures on the combinatorics of free probability, Cambridge University Press (2006).
- [69] R. Nicoara, A finiteness result for commuting squares of matrix algebras, *J. Operator Theory* **55** (2006), 295–310.
- [70] G.K. Pedersen, C^* -algebras and their automorphism groups, Academic Press (1979).
- [71] M. Petrescu, Existence of continuous families of complex Hadamard matrices of certain prime dimensions and related results, Ph.D. Thesis, UCLA (1997).
- [72] S. Popa, Orthogonal pairs of $*$ -subalgebras in finite von Neumann algebras, *J. Operator Theory* **9** (1983), 253–268.
- [73] S. Raum, Isomorphisms and fusion rules of orthogonal free quantum groups and their complexifications, *Proc. Amer. Math. Soc.* **140** (2012), 3207–3218.
- [74] P.M. Sołtan, Quantum families of maps and quantum semigroups on finite quantum spaces, *J. Geom. Phys.* **59** (2009), 354–368.
- [75] W. Tadej and K. Życzkowski, A concise guide to complex Hadamard matrices, *Open Syst. Inf. Dyn.* **13** (2006), 133–177.
- [76] W. Tadej and K. Życzkowski, Defect of a unitary matrix, *Linear Algebra Appl.* **429** (2008), 447–481.
- [77] T. Tao, Fuglede’s conjecture is false in 5 and higher dimensions, *Math. Res. Lett.* **11** (2004), 251–258.
- [78] I. Tuba and H. Wenzl, On braided tensor categories of type BCD, *J. Reine Angew. Math.* **581** (2005), 31–69.
- [79] S. Vaes and R. Vergnioux, The boundary of universal discrete quantum groups, exactness and factoriality, *Duke Math. J.* **140** (2007), 35–84.
- [80] R. Vergnioux, Orientation of quantum Cayley trees and applications, *J. Reine Angew. Math.* **580** (2005), 101–138.
- [81] D.V. Voiculescu, K.J. Dykema and A. Nica, Free random variables, AMS (1992).
- [82] C. Voigt, The Baum-Connes conjecture for free orthogonal quantum groups, *Adv. Math.* **227** (2011), 1873–1913.
- [83] S. Wang, Free products of compact quantum groups, *Comm. Math. Phys.* **167** (1995), 671–692.
- [84] S. Wang, Quantum symmetry groups of finite spaces, *Comm. Math. Phys.* **195** (1998), 195–211.
- [85] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [86] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35–76.
- [87] S.L. Woronowicz, Compact quantum groups, in “Symétries quantiques”, North-Holland (1998), 845–884.

T.B.: DEPARTMENT OF MATHEMATICS, CERGY-PONTOISE UNIVERSITY, 95000 CERGY-PONTOISE, FRANCE. teodor.banica@u-cergy.fr