

TWO-PARAMETER FAMILIES OF QUANTUM SYMMETRY GROUPS

TEODOR BANICA AND ADAM SKALSKI

ABSTRACT. We introduce and study natural two-parameter families of quantum groups motivated on one hand by the liberations of classical orthogonal groups and on the other by quantum isometry groups of the duals of the free groups. Specifically, for each pair (p, q) of non-negative integers we define and investigate quantum groups $O^+(p, q)$, $B^+(p, q)$, $S^+(p, q)$ and $H^+(p, q)$ corresponding to, respectively, orthogonal groups, bistochastic groups, symmetric groups and hyperoctahedral groups. In the first three cases the new quantum groups turn out to be related to the (dual free products of) free quantum groups studied earlier. For $H^+(p, q)$ the situation is different and we show that $H^+(p, 0) \approx \widehat{\text{QISO}(\mathbb{F}_p)}$, where the latter can be viewed as a liberation of the classical isometry group of the p -dimensional torus.

INTRODUCTION

Compact quantum groups have entered mathematics in late 1980s (see [Wor₃], [MVD] and references therein). Recent years have brought an increased interest in investigating quantum groups as quantum symmetry or isometry groups of classical or quantum spaces ([Ban₂], [BBC₁], [Gos], [BGS], [BaG]). One particular approach to constructing quantum symmetry groups is the so-called ‘liberation’ of classical compact groups. This technique, developed by the first named author and his collaborators, is based on choosing a suitable collection of relations satisfied by the functions on the group in question and then relaxing the commutativity assumptions ([BaS]). On the other hand in recent work of Bhowmick and the second named author (motivated by [Gos]) a quantum isometry group has been associated with the (dual of) each finitely generated discrete group; in particular the quantum isometry groups of the duals of the free groups were computed. The last result and the form of the obtained quantum isometry groups suggested considering a general framework in which a variation of universal quantum orthogonal groups of [VDW] is realised by replacing the usual selfadjointness of entries by imposing specific relations between entries and their adjoints. Although it turns out that this new choice actually does not introduce nontrivial modifications on the level of quantum orthogonal groups, the situation changes if we consider quantum symmetric groups ([Wan₂]) or quantum hyperoctahedral groups ([BBC₂]). In this paper we present a full study of these deformations and cast them in the language of ‘liberations’ studied in [BaS].

2000 *Mathematics Subject Classification.* Primary 46L65, Secondary 05E10, 16T05, 46L54, 46L87.

Key words and phrases. Quantum symmetry groups, quantum isometry groups, liberation, representation theory of quantum groups, Tannakian categories.

As we are going to consider at the same time deformed adjoint relations and the usual selfadjointness conditions on some other entries, we will throughout the paper work with two parameters, p (representing the deformed relations) and q (the standard ones). These parameters are assumed to be non-negative integers, not simultaneously equal to 0. As the deformed relations involve pairs of coordinates, the quantum groups that we study will have ‘rank’ (the dimension of the fundamental unitary representation) equal to $2p+q$. We will consider deformed quantum versions of orthogonal, bistochastic, symmetric and hyperoctahedral groups. Main body of the results obtained in this paper can be summarized in the following table (the categories of representations are described in terms of noncrossing, possibly ‘bulleted’, partitions, with the details given in Section 4; $i(p, q)$ is defined to be equal to 1 if $pq = 0$, and to 2 if $pq > 0$):

quantum group \mathbb{G}	$O^+(p, q)$	$B^+(p, q)$	$S^+(p, q)$	$H^+(p, q)$
algebra $C(\mathbb{G})$	$A_o(2p+q)$	$A_b(p, q)$	$A_h(p) \star A_s(q)$	$A_h(p, q)$
cat. of reps	NC_2	NC_{12}	$NC_\bullet \star NC$	NC_{even}
classical version	O_{2p+q}	$O_{2p+q-i(p,q)}$	$H_p \times S_q$	$(\mathbb{T}^p \rtimes H_p) \times H_q$

As explained earlier, if $p = 0$ we obtain the ‘liberated objects’ studied in [BaS]. On the other hand when $q = 0$ we obtain the following (in particular $H^+(p, 0)$ is the quantum isometry group of the C^* -algebra of the free group \mathbb{F}_p discovered in [BhS] and in a way providing a starting point for the considerations in this work):

quantum group \mathbb{G}	$O^+(p, 0)$	$B^+(p, 0)$	$S^+(p, 0)$	$H^+(p, 0)$
quantum symmetry group of	\mathbb{S}^{2p}	\mathbb{S}_-^{2p}	$[0, 1]^p$	$\widehat{\mathbb{F}}_p$
classical version	O_{2p}	O_{2p-1}	H_p	$\mathbb{T}^p \rtimes H_p$
classical symmetry group of	\mathcal{S}^{2p}	\mathcal{S}_-^{2p}	$[0, 1]^p$	\mathbb{T}^p

Above \mathcal{S}^{2p} and \mathbb{S}^{2p} denote respectively the usual sphere in \mathbb{R}^{2p} and the free sphere studied in [BaG], and \mathcal{S}_-^{2p} and \mathbb{S}_-^{2p} denote the respective spheres with one coordinate fixed. With the isomorphisms established in this paper the first two columns can be deduced from results in [BaG] (and the third is a consequence of [BBC₂]).

The detailed plan of the paper is as follows: in Section 1 we quote basic definitions, establish some terminology related to compact quantum groups and recall the quantum (free) symmetry groups corresponding to orthogonal, symmetric, bistochastic and hyperoctahedral groups. In Section 2 the study of their two-parameter counterparts begins with the analysis of the orthogonal, bistochastic and symmetric quantum groups denoted respectively $O^+(p, q)$, $B^+(p, q)$ and $S^+(p, q)$. It turns out that all of them can be described in terms of (the free products of) the one-parameter versions. Section 3 contains a detailed analysis of the two-parameter quantum hyperoctahedral group $H^+(p, q)$; in particular we show that $H^+(p, 0)$ coincides with the quantum isometry group of the dual of the free group discovered in [BhS]. Section 4 is devoted to establishing the description of the categories of representations of our quantum groups in terms of non-crossing (marked) partitions. Further in Section 5 theorems proved in Sections 2-4 are used to analyse the relations between the quantum groups studied in the paper: we investigate the generation results, intersection results and inclusions of the form $\mathbb{G}(p, 0) \hat{\star} \mathbb{G}(0, q) \subset \mathbb{G}(p, q)$.

We also show there a fact conjectured in [BlS]: the two-parameter quantum hyperoctahedral group $H^+(p, q)$ may be viewed as a quantum extension of the quantum symmetric group S_{2p+q}^+ . In Section 6 we describe the classical versions of the quantum groups we study to show that each of these quantum groups has a natural description as a liberation of a classical symmetry group. Finally in the last section we discuss in what sense the family of quantum groups described in this paper exhausts the natural two-parameter construction presented in Sections 2-3 and in the process discover another two-parameter quantum group, $H_s^+(p, q)$ which turns out to be isomorphic to the dual free product of H_p^{+4} and H_q^+ .

1. COMPACT QUANTUM GROUPS - DEFINITIONS AND NOTATION

In this section we recall the definition of a compact quantum group due to Woronowicz and introduce quantum orthogonal, symmetric, hyperoctahedral and bistochastic groups. The minimal tensor product of C^* -algebras will be denoted by \otimes , algebraic tensor products by \odot . For $n \in \mathbb{N}$ we denote the algebra of n by n complex matrices by M_n .

Definition 1.1. Let A be a unital C^* -algebra and $\Delta : A \rightarrow A \otimes A$ be a unital $*$ -homomorphism satisfying the coassociativity condition:

$$(\Delta \otimes \text{id}_A)\Delta = (\text{id}_A \otimes \Delta)\Delta.$$

If additionally $\overline{\Delta(A)(1 \otimes A)} = \overline{\Delta(A)(A \otimes 1)} = A \otimes A$ we say that A is the algebra of continuous functions on a compact quantum group \mathbb{G} and usually write $A = C(\mathbb{G})$. A unique dense Hopf $*$ -subalgebra of $C(\mathbb{G})$, the algebra of coefficients of finite-dimensional unitary representations of \mathbb{G} , will be denoted by $R(\mathbb{G})$.

If \mathbb{G} is a compact quantum group and $n \in \mathbb{N}$ then a unitary matrix $U = (U_{ij})_{i,j=1}^n \in M_n(C(\mathbb{G}))$ is called a *fundamental representation* of \mathbb{G} (or a *fundamental corepresentation* of $C(\mathbb{G})$) if for each $i, j = 1, \dots, n$

$$\Delta(U_{ij}) = \sum_{k=1}^n U_{ik} \otimes U_{kj}$$

and the entries of U generate $C(\mathbb{G})$ as a C^* -algebra. If \mathbb{G} admits a fundamental representation, it is called a *compact matrix quantum group*. This will be the case for all quantum groups considered in this paper; in fact they will be defined via their respective fundamental representations.

In general there is an ambiguity in passing from $R(\mathbb{G})$ to $C(\mathbb{G})$ related to the fact that not all compact quantum groups are *coamenable*. As all quantum groups studied in this paper will be defined by universal properties, we will assume that $C(\mathbb{G})$ is the *universal* completion of $R(\mathbb{G})$ ([BMT]).

We will later need a free product construction introduced in [Wan₁]. If $\mathbb{G}_1, \mathbb{G}_2$ are compact quantum groups, then the C^* -algebraic free product $C(\mathbb{G}_1) \star C(\mathbb{G}_2)$ has a natural structure of the algebra of continuous functions on a compact quantum group, to be denoted $\mathbb{G}_1 \hat{\star} \mathbb{G}_2$. In particular if $U_1 \in M_n(C(\mathbb{G}_1))$ and $U_2 \in M_n(C(\mathbb{G}_2))$ are respective fundamental corepresentations, then $\begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix} \in M_n(C(\mathbb{G}_1) \star C(\mathbb{G}_2))$ is the fundamental corepresentation of $C(\mathbb{G}_1 \hat{\star} \mathbb{G}_2)$. The construction is dual to the usual free product of discrete groups: when the quantum

groups in question are duals of classical discrete groups, $\mathbb{G}_1 = \widehat{\Gamma}_1$, $\mathbb{G}_2 = \widehat{\Gamma}_2$, then $\mathbb{G}_1 \widehat{\star} \mathbb{G}_2 \approx \widehat{\Gamma_1 \star \Gamma_2}$.

Let $n \in \mathbb{N}$; it will denote the dimension of the fundamental representation of the compact quantum groups defined below. The following definition comes from [VDW]; here we recast it in the language described above.

Definition 1.2. Let $F \in M_n$ be an invertible matrix such that $F\bar{F} = cI_n$ for some $c \in \mathbb{C}$. Let $A_o(F)$ denote the universal C^* -algebra generated by the entries of a unitary $U \in M_n \otimes A_o(F)$ such that

$$(1.1) \quad U = (F \otimes 1)\bar{U}(F^{-1} \otimes 1)$$

(here and in what follows a bar over the matrix denotes a matrix obtained by an entrywise conjugation of entries). When $U \in M_n \otimes A_o(F)$ is interpreted as the fundamental unitary corepresentation, we can view $A_o(F)$ as the algebra of continuous functions on the compact quantum group denoted by $O^+(F)$. In particular if $F = I_n$, we write simply $A_o(n)$ and O_n^+ instead of $A_o(I_n)$ and $O^+(I_n)$.

Recall that if \mathbf{A} is a C^* -algebra then a unitary matrix $U \in M_n(\mathbf{A})$ is called a *magic unitary* if each entry of U is a projection. A unitary $U \in M_n(\mathbf{A})$ is called *cubic* if its entries are selfadjoint and the products of different entries lying in the same row or column are 0. The following definitions come respectively from [Wan₂] and [BBC₂].

Definition 1.3. Denote by $A_s(n)$ the universal C^* -algebra generated by the entries of an n by n magic unitary U . When $U \in M_n \otimes A_s(n)$ is interpreted as the fundamental unitary corepresentation, we view $A_s(n)$ as the algebra of continuous functions on the quantum permutation group on n elements, S_n^+ .

Definition 1.4. Denote by $A_h(n)$ the universal C^* -algebra generated by the entries of an n by n cubic unitary U . When $U \in M_n \otimes A_h(n)$ is interpreted as the fundamental unitary corepresentation, we view $A_h(n)$ as the algebra of continuous functions on the quantum hyperoctahedral group on n coordinates, H_n^+ .

The quantum groups O_n^+ , S_n^+ and H_n^+ are also called the free orthogonal quantum group, the free symmetric quantum group and the free hyperoctahedral quantum group and can be respectively viewed as liberations of the compact groups O_n , S_n and H_n ([BaS]). More information about their properties, including their interpretations as quantum symmetry groups can be found in that paper and also respectively in [BaG], [BBC₁] and [BBC₂].

The following definition was introduced in [BaS].

Definition 1.5. Let $n \in \mathbb{N}$. Denote by $A_b(n)$ the universal C^* -algebra generated by the entries of an n by n unitary U , which has selfadjoint entries which sum to 1 in each row/column. When $U \in M_n \otimes A_b(n)$ is interpreted as the fundamental unitary corepresentation, we view $A_b(n)$ as the algebra of continuous functions on the quantum bistochastic group in n dimensions, B_n^+ .

The condition on the sum of entries in each row/column being equal to 1 is equivalent to stating that the vector $[1, \dots, 1]^t$ is an eigenvector for both U and U^t . This observation leads to a natural isomorphism $B_n^+ \approx O_{n-1}^+$ ([Rau]).

Relations between the C^* -algebras and quantum groups introduced above can be expressed via the following diagrams (arrows on the level of algebras denote

surjective unital *-homomorphisms intertwining the respective coproducts):

$$(1.2) \quad \begin{array}{ccc} A_o(n) & \rightarrow & A_b(n) & & O_n^+ & \supset & B_n^+ \\ & & \downarrow & & \downarrow & & \cup & & \cup \\ & & A_h(n) & \rightarrow & A_s(n) & & H_n^+ & \supset & S_n^+ \end{array}$$

The diagram on the right suggests a number of questions, for instance whether $O_n^+ = \langle B_n^+, H_n^+ \rangle$, or whether $S_n^+ = B_n^+ \cap H_n^+$ (at the level of classical versions, the answers are yes and yes). Once these questions are properly formulated, for instance in terms of tensor categories, the answers turn out to be positive, and can be deduced from [BaS]. We will describe the details later, when we discuss similar problems in a more general 2-parameter framework.

2. QUANTUM GROUPS $O^+(p, q)$, $B^+(p, q)$ AND $S^+(p, q)$

In this section we describe the two-parameter families of quantum groups generalising the quantum orthogonal, bistochastic and symmetric groups described in Section 1. We begin by introducing necessary notations.

Let $p, q \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, $p^2 + q^2 > 0$ and let $\rho = e^{\frac{2\pi i}{8}}$. Put $\mathcal{F} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\mathcal{C} = \frac{1}{\sqrt{2}} \begin{pmatrix} \rho & \rho^7 \\ \rho^3 & \rho^5 \end{pmatrix}$, let $F_{p,0} \in M_{2p}$ be the matrix given by

$$F_{p,0} = \begin{pmatrix} \mathcal{F} & 0 & \cdots & 0 \\ 0 & \mathcal{F} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \mathcal{F} \end{pmatrix}$$

and define $F_{p,q} \in M_{2p+q}$ by

$$(2.1) \quad F_{p,q} = \begin{pmatrix} F_{p,0} & 0 \\ 0 & I_q \end{pmatrix}$$

The matrix $F_{p,q}$ is a selfadjoint unitary, and a permutation matrix; moreover $\overline{F_{p,q}} = (F_{p,q})^t = (F_{p,q})^*$. The matrix $C_{p,q} \in M_{2p+q}$ is defined in an analogous way, replacing \mathcal{F} by \mathcal{C} in appropriate matrix blocks. Note that \mathcal{C} , so also $C_{p,q}$, is unitary.

Whenever we consider matrices of the size $2p + q$ we will denote the indices corresponding to the ‘ p -part’ by pairs $i\alpha$, where $i \in \{0, 1\}$ and $\alpha \in \{1, \dots, p\}$ and to the ‘ q -part’ by capital Latin letters running from 1 to q . Moreover we will use the notation \bar{i} (where $\bar{0} = 1$, $\bar{1} = 0$, so that $\bar{\bar{i}} = i$). This facilitates the description of the fact that coordinates in the ‘ p -part’ come naturally in pairs. To simplify the notation we will also write $\mathcal{J}_p = \{i\alpha : i \in \{0, 1\}, \alpha \in \{1, \dots, p\}\}$, $\mathcal{J}_{p,q} = \mathcal{J}_p \cup \{1, \dots, q\}$, and extend the ‘bar’ notation to indices in $\mathcal{J}_{p,q}$, writing $\bar{z} = \bar{i}\alpha$ if $z = i\alpha \in \mathcal{J}_p$ and $\bar{z} = M$ if $z = M \in \{1, \dots, q\}$. The canonical basis in \mathbb{C}^{2p+q} will be often denoted by $(e_z)_{z \in \mathcal{J}_{p,q}}$.

2.1. Quantum group $O^+(p, q) \approx O_{2p+q}^+$. As the matrix $F_{p,q}$ defined in (2.1) satisfies the conditions listed in Definition 1.2 we can consider $A_o(p, q) := A_o(F_{p,q})$ and $O^+(p, q) := O^+(F_{p,q})$. Clearly $O^+(0, q) \approx O_q^+$; in fact the discussion in Section

5 of [BDV] implies that $O^+(p, q) \approx O_{2p+q}^+$. From our point of view it is important to consider the following rephrasing of the above definitions:

Theorem 2.1. *The algebra $A_o(p, q)$ is the universal C^* -algebra generated by elements $\{U_{z,y} : z, y \in \mathcal{J}_{p,q}\}$ such that the resulting $2p + q$ by $2p + q$ matrix U is unitary and for each $i\alpha, j\beta \in \mathcal{J}_p, M, N \in \{1, \dots, q\}$ we have*

$$(2.2) \quad U_{i\alpha, j\beta}^* = U_{i\alpha, \bar{j}\beta},$$

$$(2.3) \quad U_{i\alpha, N}^* = U_{i\alpha, N},$$

$$(2.4) \quad U_{M, j\beta}^* = U_{M, \bar{j}\beta},$$

$$(2.5) \quad U_{M, N}^* = U_{M, N}.$$

Moreover $A_o(p, q)$ with U viewed as a fundamental corepresentation is the algebra of continuous functions on a compact quantum group $O^+(p, q) \approx O_{2p+q}^+$.

Proof. The first part is a direct consequence of the fact that $F_{p,q}$ has a block-matrix form and formulas:

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} A^* & B^* \\ C^* & D^* \end{bmatrix} = \begin{bmatrix} C^* & D^* \\ A^* & B^* \end{bmatrix}, \quad \begin{bmatrix} A^* & B^* \\ C^* & D^* \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} B^* & A^* \\ D^* & C^* \end{bmatrix}$$

and

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} A^* & B^* \\ C^* & D^* \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} D^* & C^* \\ B^* & A^* \end{bmatrix},$$

which imply that if $U = (U_{z,y})_{z,y \in \mathcal{J}_{p,q}}$ then $U = (F_{p,q} \otimes 1)\bar{U}(F_{p,q} \otimes 1)$ if and only if the entries of U satisfy relations (2.2)-(2.5).

The second part follows from the discussion before the theorem, but can be also seen directly: indeed, exploiting the equalities of the type

$$\begin{bmatrix} \rho & \rho^7 \\ \rho^3 & \rho^5 \end{bmatrix} \begin{bmatrix} A \\ C \end{bmatrix} = \begin{bmatrix} \rho(A + \rho^6 C) \\ \rho^3(A + \rho^2 C) \end{bmatrix},$$

$$\begin{bmatrix} A & B \end{bmatrix} \begin{bmatrix} \rho^7 & \rho^5 \\ \rho & \rho^3 \end{bmatrix} = [\rho(\rho^6 A + B) \quad \rho^3(\rho^2 A + B)]$$

and

$$\begin{bmatrix} \rho & \rho^7 \\ \rho^3 & \rho^5 \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} \rho^7 & \rho^5 \\ \rho & \rho^3 \end{bmatrix} = \begin{bmatrix} A - iC + iB + D & -iA - C - B + iD \\ iA - C - B - iD & A + iC - iB + D \end{bmatrix}$$

one can check that if A is a C^* -algebra and $U \in M_{2p+q}(A)$ is a unitary matrix, then the entries of U satisfy conditions displayed in the theorem if and only if the entries of the unitary matrix $(C_{p,q} \otimes 1_A)U(C_{p,q}^* \otimes 1_A)$ are self-adjoint. \square

Note that in terms of the notation introduced earlier the relations (2.2)-(2.5) can be summarized by saying

$$(2.6) \quad U_{z,y} = U_{\bar{z}, \bar{y}}^*, \quad z, y \in \mathcal{J}_{p,q}.$$

It is well known that algebraic relations between entries of the fundamental representation U of a given quantum group can be interpreted as declaring certain scalar matrices to be elements of $\text{Hom}(U^{\otimes k}; U^{\otimes l})$ for some $k, l \in \mathbb{N}$ (see for example [Ban₁]).

Proposition 2.2. *The algebra $A_o(p, q)$ is the universal C^* -algebra generated by the entries of a unitary $2p+q$ by $2p+q$ matrix U such that the vector $\xi := \sum_{i\alpha \in \mathcal{J}_p} e_{i\alpha} \otimes e_{\bar{i}\alpha} + \sum_{M=1}^q e_M \otimes e_M$ is a fixed vector for $U^{\otimes 2}$ (in other words the map $1 \rightarrow \xi$ is an element of $\text{Hom}(1; U^{\otimes 2})$).*

Proof. Let $(U_{z,y})_{z,y \in \mathcal{J}_{p,q}}$ be unitary. Then $U^{\otimes 2}\xi = \xi$ if and only if

$$\sum_{z,y \in \mathcal{J}_{p,q}} \left(\sum_{i\alpha \in \mathcal{J}_p} U_{z,i\alpha} U_{y,\bar{i}\alpha} + \sum_{M=1}^q U_{z,M} U_{y,M} \right) e_z \otimes e_y = \xi.$$

This means that for example for each $j\beta \in \mathcal{J}_q$ and $y \in \mathcal{J}_{p,q}$ we must have

$$\sum_{i\alpha \in \mathcal{J}_p} U_{j\beta,i\alpha} U_{y,\bar{i}\alpha} + \sum_{M=1}^q U_{j\beta,M} U_{y,M} = \delta_y^{\bar{j}\beta}.$$

Fix $z \in \mathcal{J}_{p,q}$, multiply the last formula by $U_{y,z}^*$ on the right and sum over $y \in \mathcal{J}_{p,q}$. As U is unitary, this yields

$$\sum_{i\alpha \in \mathcal{J}_p} U_{j\beta,i\alpha} \delta_{i\alpha}^z + \sum_{M=1}^q U_{j\beta,M} \delta_M^z = U_{j\beta,z}^*.$$

Considering respectively $z \in \{1, \dots, q\}$ and $z \in \mathcal{J}_p$ yields the first two displayed formulas in Theorem 2.1. It is easy to see that the third and fourth formula can be obtained in an analogous way. \square

The last proposition allows one to describe the tensor category corresponding to the dual of $O^+(p, q)$ in terms of Temperley-Lieb diagrams ([Ban₁]) or noncrossing pair partitions ([BaS]). We refer to these papers for details; in Section 4 below we state the corresponding results in terms of partitions.

2.2. Quantum group $B^+(p, q) \approx O_{p+q-i(p,q)}^+$

Definition 2.3. The algebra $A_b(p, q)$ is the universal C^* -algebra generated by elements $\{U_{z,y}, z, y \in \mathcal{J}_{p,q}\}$ which satisfy all the relations required of generators of $A_o(p, q)$ and additionally are such that entries in each row/column of the resulting unitary U sum to 1: for all $y \in \mathcal{J}_{p,q}$

$$\sum_{z \in \mathcal{J}_{p,q}} U_{z,y} = \sum_{z \in \mathcal{J}_{p,q}} U_{y,z} = 1.$$

Let $\eta = \sum_{z \in \mathcal{J}_{p,q}} e_z \in \mathbb{C}^{2p+q}$ and recall the matrix $C_{p,q}$ defined in the beginning of this section.

Theorem 2.4. *The algebra $A_b(p, q)$ is the algebra of continuous functions on a compact quantum group, denoted further $B^+(p, q)$. The unitary $U = (U_{z,y})_{z,y \in \mathcal{J}_{p,q}} \in M_{2p+q} \otimes A_b(p, q)$ is the fundamental representation of $B^+(p, q)$. The algebra $A_b(p, q)$ is isomorphic to the universal C^* -algebra generated by entries of a $2p+q$ by $2p+q$ unitary V which is orthogonal and satisfies the condition $V(C_{p,q}\eta \otimes 1) = C_{p,q}\eta \otimes 1$.*

Proof. As explained after Definition 1.5 the condition that the entries in each row and column of a matrix U sum to 1 are equivalent to the fact that the vector η is fixed both by U and U^t . This observation (or a direct computation) implies that $A_b(p, q)$ is the algebra of continuous functions on a compact quantum group. Further note that as $F_{p,q}\eta = \eta$ and $F_{p,q}$ is a selfadjoint unitary we have the following

string of equivalences for a unitary $U \in M_n(A_o(p, q))$ satisfying the condition (1.1) with $F = F_{p,q}$:

$$\begin{aligned} U(\eta \otimes 1) = \eta \otimes 1 &\Leftrightarrow U^*(\eta \otimes 1) = \eta \otimes 1 \Leftrightarrow (F_{p,q} \otimes 1)U^t(F_{p,q} \otimes 1)(\eta \otimes 1) = \eta \otimes 1 \\ &\Leftrightarrow U^t(\eta \otimes 1) = \eta \otimes 1, \end{aligned}$$

so that the condition on the sum of entries in each column of a unitary U as above being equal to 1 follows from the analogous condition for rows.

In the last part of the proof of Theorem 2.1 we noticed that the transformation between the fundamental unitary in $O^+(p, q)$ and that of O_{2p+q}^+ is implemented by conjugating with the unitary matrix $C_{p,q}$. Hence to prove the last statement it suffices to note that a unitary U fixes the vector η if and only if $(C_{p,q} \otimes 1)U(C_{p,q}^* \otimes 1)$ fixes $C_{p,q}\eta$. \square

When $q = 0$ or $p = 0$ the above theorem implies that the ‘deformed’ quantum bistochastic group coincides with the one studied in [BaS].

Theorem 2.5. *The following isomorphisms hold:*

$$\begin{aligned} B^+(p, 0) &\approx B_{2p}^+ \approx O_{2p-1}^+, & B^+(0, q) &\approx B_q^+ \approx O_{q-1}^+, \\ B^+(p, q) &\approx O_{2p+q-2}^+ \text{ for } pq > 0, \\ B^+(p, 1) &\approx B^+(p, 0). \end{aligned}$$

Proof. The only isomorphism in the first line that needs to be proved is $B^+(p, 0) \approx B_{2p}^+$ (the ones involving the quantum orthogonal groups are the consequences of [Rau], as explained after Definition 1.5). Due to Theorem 2.4 the fundamental representation of $B^+(p, 0)$ can be defined as a $2p$ by $2p$ unitary matrix U with selfadjoint entries which satisfies the condition $U(C_{p,0}\eta \otimes 1) = C_{p,0}\eta \otimes 1$. It is clear that in the above condition $C_{p,0}\eta$ can be replaced by a non-zero scalar multiple; in particular by the vector $[1, -1, \dots, 1, -1]^t$. But that vector can be mapped by a real orthogonal matrix onto η , so also onto $[1, 0, 0, \dots, 0]^t$, and the argument of [Rau] implies that the desired isomorphism holds.

Assume now that $pq > 0$. Using once again Theorem 2.4 a fundamental representation of $B^+(p, q)$ can be defined as a $2p + q$ by $2p + q$ unitary matrix U with selfadjoint entries which satisfies the condition $U(C_{p,q}\eta \otimes 1) = C_{p,q}\eta \otimes 1$. One can check that the vector $C_{p,q}\eta$ is proportional to the vector $\tilde{\eta}_{p,q} \in \mathbb{C}^{2p+q}$ given by

$$\begin{aligned} \tilde{\eta}_{p,q} &= \underbrace{[1 - i, i - 1, \dots, 1 - i, i - 1]}_{2p \text{ times}} \underbrace{[1 + i, \dots, 1 + i]}_{q \text{ times}}^T \\ &= \underbrace{[1, -1, \dots, 1, -1]}_{2p \text{ times}} \underbrace{[1, \dots, 1]}_{q \text{ times}}^T + i \underbrace{[-1, 1, \dots, -1, 1]}_{2p \text{ times}} \underbrace{[1, \dots, 1]}_{q \text{ times}}^T \end{aligned}$$

As U has selfadjoint entries, it preserves $\tilde{\eta}_{p,q}$ if and only if it preserves its real and imaginary parts; equivalently, it preserves vectors

$$\underbrace{[1, -1, \dots, 1, -1]}_{2p \text{ times}} \underbrace{[0, \dots, 0]}_{q \text{ times}}^T$$

and

$$\underbrace{[0, 0, \dots, 0, 0]}_{2p \text{ times}} \underbrace{[1, \dots, 1]}_{q \text{ times}}^T.$$

Repeating an earlier argument we find a matrix in $O_{2p} \times O_q \subset O_{2p+q}$ mapping these vectors respectively to $[1, 0, \dots, 0]^T$ and $[0, \dots, 0, 1]^T$; this provides the isomorphism $B^+(p, q) \approx O_{2p+q-2}^+$, which implies in particular that $B^+(p, 1) \approx B_{p,0}^+$. \square

Note that the argument in the second part of the above proof can be framed in general terms – for $n \geq 3$ the group of n by n orthogonal matrices preserving a fixed non-zero vector $v \in \mathbb{C}^n$ is either isomorphic to O_{n-1} (when the real and imaginary parts of v are proportional to each other) or to O_{n-2} (when they are not); the same applies to O_n^+ .

The following result is a consequence of Proposition 2.2 and the arguments in the proof of Theorem 2.4.

Corollary 2.6. *The algebra $A_b(p, q)$ is the universal C^* -algebra generated by the entries of a unitary $2p + q$ by $2p + q$ matrix U such that the vector ξ defined in Proposition 2.2 is a fixed vector for $U^{\otimes 2}$ and the vector $\eta = \sum_{z \in \mathcal{J}_{p,q}} e_z$ is a fixed vector for U .*

The above corollary implies that the category of representations of $B^+(p, q)$ coincides with that of $B^+(2p + q)$ ([BaS]), see Section 4.

2.3. Quantum group $S^+(p, q) \approx H_p^+ \hat{\star} S_q^+$. The free quantum group of permutations of n elements may be viewed as the universal C^* -algebra generated by entries of an orthogonal matrix which are additionally required to be orthogonal projections. This motivates the following definition.

Definition 2.7. The algebra $A_s(p, q)$ is the universal C^* -algebra generated by projections $\{U_{z,y}, z, y \in \mathcal{J}_{p,q}\}$ which satisfy all the relations required of generators of $A_o(p, q)$.

As for every magic unitary entries lying in the same row or column are pairwise orthogonal, the generators of $A_s(p, q)$ satisfy the following relations:

$$U_{z,y}^* U_{z,x} = U_{z,y} U_{z,x} = 0, \quad U_{y,z}^* U_{x,z} = U_{y,z} U_{x,z} = 0 \quad z, y, x \in \mathcal{J}_{p+q}, x \neq y.$$

Proposition 2.8. *The algebra $A_s(p, q)$ is the universal C^* -algebra generated by two families of projections $\{U_{i\alpha, j\beta} : i\alpha, j\beta \in \mathcal{J}_p\}$ and $\{U_{M,N} : M, N \in \{1, \dots, q\}\}$, such that both matrices $(U_{i\alpha, j\beta})_{i\alpha, j\beta \in \mathcal{J}_p}$ and $(U_{M,N})_{M,N=1}^q$ are magic unitaries and moreover*

$$(2.7) \quad U_{i\alpha, j\beta} = U_{i\alpha, \bar{j}\beta}, \quad i\alpha, j\beta \in \mathcal{J}_p.$$

Proof. It suffices to show that whenever $i\alpha \in \mathcal{J}_p$ and $M \in \{1, \dots, q\}$ then $U_{i\alpha, M} = 0 = U_{M, i\alpha}$. But the matrix $\{U_{z,y}, z, y \in \mathcal{J}_{p,q}\} \in M_{2p+q}(A_s(p, q))$ is a magic unitary, so each of its columns consists of mutually orthogonal projections, and as we have $U_{i\alpha, M} = U_{i\alpha, M}^*$, it follows that $U_{i\alpha, M} = 0$. The second equality follows from the orthogonality of projections in each row of a magic unitary. \square

Theorem 2.9. *The algebra $A_s(p, q)$ is the algebra of continuous functions on a compact quantum group, denoted further $S^+(p, q)$. The unitary $U = (U_{z,y})_{z,y \in \mathcal{J}_{p,q}} \in M_{2p+q} \otimes A_s(p, q)$ is the fundamental representation of $S^+(p, q)$. The algebra $A_s(p, q)$ is isomorphic to the free product $A_h(p) \star A_s(q)$; on the level of quantum groups we have $S^+(p, q) \approx H_p^+ \hat{\star} S_q^+$.*

Proof. Due to Proposition 2.8 it suffices to consider separately the cases $p = 0$ and $q = 0$; as we have $A_s(0, q) \approx A_s(q)$ (with the isomorphism preserving natural fundamental corepresentations), we can assume that $q = 0$. Further it suffices to show that if we define for each $i\alpha, j\beta \in \mathcal{J}_p$

$$\tilde{U}_{i\alpha, j\beta} := \sum_{k\gamma \in \mathcal{J}_p} U_{i\alpha, k\gamma} \otimes U_{k\gamma, j\beta}$$

then each $\tilde{U}_{i\alpha, j\beta}$ is a projection (the fact that the conditions in (2.7) will then be satisfied follows from Theorem 2.1; and analogous statements for the potential antipode and counit follow from the fact that the adjoint of a projection is a projection and $0, 1 \in \mathbb{C}$ are projections). The last statement is however a direct consequence of orthogonality of the rows/columns of the magic unitary; it can also be deduced from the fact that the map $U_{i\alpha, j\beta} \rightarrow \tilde{U}_{i\alpha, j\beta}$ is a $*$ -homomorphism.

Proposition 2.8 implies also that $A_s(p, q) \approx A_s(p, 0) \star A_s(0, q)$. Therefore it suffices to show that $A_s(p, 0) \approx A_h(p)$. This is however an immediate consequence of the fact that $A_h(p)$ can be defined via the requirement that its fundamental corepresentation is a $2p$ by $2p$ *sudoku* unitary, i.e. a magic unitary which has a block matrix form $\begin{pmatrix} a & b \\ b & a \end{pmatrix}$ (Definition 5.2 and Theorem 6.2 of [BBC₂]). Indeed, formulas (2.2)-(2.5) imply that the fundamental unitary representation of $S^+(p, 0)$ is a magic unitary of the form

$$\begin{bmatrix} A & B & C & D & E & F & \dots \\ B & A & D & C & F & E & \dots \\ G & H & I & J & K & L & \dots \\ H & G & J & I & L & K & \dots \\ M & N & O & P & Q & R & \dots \\ N & M & P & O & R & Q & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

which can be transformed into a sudoku unitary by permuting rows and columns (so that odd rows/columns remain in the same order but are shifted to the left/up so that they become first p rows/columns). \square

The next proposition facilitates the description of the category of the representations of $S^+(p, q)$ in terms of the noncrossing partitions, see Section 4.

Proposition 2.10. *The algebra $A_s(p, q)$ is the universal C^* -algebra generated by the entries of a unitary $2p + q$ by $2p + q$ matrix U such that the vector ξ defined in Proposition 2.2 is a fixed vector for $U^{\otimes 2}$ and the map $e_{i\alpha} \rightarrow e_{i\alpha} \otimes e_{\bar{i\alpha}}$, $e_M \rightarrow e_M \otimes e_M$ defines a morphism in $\text{Hom}(U; U^{\otimes 2})$.*

Proof. If U is a unitary $2p + q$ by $2p + q$ matrix such that ξ is a fixed vector for $U^{\otimes 2}$, then by Proposition 2.2 the entries of U satisfy the relations (2.2)-(2.5). Further the condition that the map described in this proposition is an intertwiner between U and $U^{\otimes 2}$ is satisfied if and only if for each $i\alpha \in \mathcal{J}_p$

$$\sum_{j\beta \in \mathcal{J}_p} e_{j\beta} \otimes e_{\bar{j\beta}} \otimes U_{j\beta, i\alpha} + \sum_{N=1}^q e_N \otimes e_N \otimes U_{N, i\alpha} = \sum_{z, y \in \mathcal{J}_{p+q}} e_z \otimes e_y \otimes U_{z, i\alpha} U_{y, \bar{i\alpha}}$$

and for each $M \in \{1, \dots, q\}$

$$\sum_{j\beta \in \mathcal{J}_p} e_{j\beta} \otimes e_{\bar{j}\beta} \otimes U_{j\beta, M} + \sum_{N=1}^q e_N \otimes e_N \otimes U_{N, M} = \sum_{z, y \in \mathcal{J}_{p+q}} e_z \otimes e_y \otimes U_{z, M} U_{y, M}.$$

The above conditions hold if and only if for each $i\alpha, j\beta \in \mathcal{J}_p, M, N \in \{1, \dots, q\}, y \in \mathcal{J}_{p+q}$

$$U_{j\beta, i\alpha} U_{y, \bar{i}\alpha} = \delta_y^{\bar{j}\beta} U_{j\beta, i\alpha},$$

$$U_{N, i\alpha} U_{y, \bar{i}\alpha} = \delta_y^N U_{N, i\alpha},$$

$$U_{j\beta, M} U_{y, M} = \delta_y^{\bar{j}\beta} U_{j\beta, M},$$

$$U_{N, M} U_{y, M} = \delta_y^N U_{N, M}.$$

Bearing in mind the relations (2.2)-(2.5) we see that all entries of U are orthogonal projections; hence U is a magic unitary. Conversely it is easy to check that if U is a magic unitary whose entries satisfy (2.2)-(2.5) then the last four displayed formulas automatically hold (see also the comment after Definition 2.7). \square

3. QUANTUM GROUP $H^+(p, q)$

The quantum groups we defined in the last section have all been shown to be closely related to well-studied objects. The generalized quantum hyperoctahedral groups to be introduced here are genuinely new quantum groups, connected to quantum hyperoctahedral groups ([BBC₂]) and quantum isometry groups of C^* -algebras of free groups ([BhS]).

The fact that an element x of a C^* -algebra is an orthogonal projection can be written as $x = x^*x$. It is natural to consider the condition $x = xx^*x$, which of course means that x is a partial isometry. This leads to the following definition.

Definition 3.1. The algebra $A_h(p, q)$ is the universal C^* -algebra generated by partial isometries $\{U_{z, y} : z, y \in \mathcal{J}_{p+q}\}$ which satisfy all the relations required of generators of $A_o(p, q)$.

Here we also have some automatic ‘orthogonality’, which will be described by the next proposition and its corollary.

Proposition 3.2. *Let A be a C^* -algebra, $n \in \mathbb{N}$ and let $U \in M_n(A)$ be a unitary matrix whose entries are partial isometries. Then*

$$U_{y, z} U_{x, z}^* = U_{z, y}^* U_{z, x} = 0, \quad z, y, x \in \{1, \dots, n\}, x \neq y.$$

Proof. For each $y, z \in \{1, \dots, n\}$ denote the initial and range projections of $U_{y, z}$ respectively by $P_{y, z}$ and $Q_{y, z}$, so that

$$P_{y, z} = U_{y, z}^* U_{y, z}, \quad Q_{y, z} = U_{y, z} U_{y, z}^*.$$

Fix $z \in \{1, \dots, n\}$. The unitarity of U implies that $\sum_{y=1}^n Q_{z, y} = \sum_{y=1}^n P_{y, z} = 1_A$, so that for $y, x \in \{1, \dots, n\}, x \neq y$ there is $Q_{z, y} Q_{z, x} = P_{y, z} P_{x, z} = 0$. The initial/range projection interpretation ends the proof. \square

Application of the above proposition and the equality (2.6) immediately gives the following corollary.

Corollary 3.3. *Let $U \in M_{2p+q}(A_h(p, q))$ be the unitary matrix ‘generating’ $A_h(p, q)$. Then*

$$U_{y,z}U_{x,z}^* = U_{z,y}^*U_{z,x} = U_{y,z}^*U_{x,z} = U_{z,y}U_{z,x}^* = 0, \quad z, y, x \in \mathcal{J}_{p,q}, x \neq y.$$

The last observations lead to the following theorem.

Theorem 3.4. *The algebra $A_h(p, q)$ is the algebra of continuous functions on a compact quantum group, denoted further $H^+(p, q)$. The unitary $U = (U_{z,y})_{z,y \in \mathcal{J}_{p,q}} \in M_{2p+q} \otimes A_h(p, q)$ is the fundamental representation of $H^+(p, q)$. The quantum group $H^+(0, q)$ is the quantum hyperoctahedral group H_q^+ studied in [BBC₂].*

Proof. The first statement can be deduced as in the proof of Theorem 2.7, using the fact that a *-homomorphic image (and the adjoint) of a partial isometry is a partial isometry. It remains to observe that the algebra $A_h(0, q)$ is isomorphic to $A_h(q)$. But this is a natural consequence of Corollary 3.3 and the fact that if a unitary is cubic then its entries satisfy the condition $u_{ij} = u_{ij}^3 = u_{ij}^*$ (easy to show and noted implicitly in [BBC₂]). \square

As stated in the introduction, the quantum group $H^+(p, 0)$ is in fact the quantum isometry group of the dual of the free group \mathbb{F}_p ([BhS]). Let us quickly recall the general notion of a quantum isometry group of the dual of a finitely generated discrete group.

Let Γ be a finitely generated discrete group with a fixed finite symmetric set of generators $S \subset \Gamma$. The choice of a generating set S determines a word-length function l on Γ . Denote the universal group C^* -algebra of Γ by $C(\widehat{\Gamma})$ and let the group ring $\mathbb{C}[\Gamma] \subset C(\widehat{\Gamma})$ be denoted by $R(\widehat{\Gamma})$. Then the multiplication by the length function defines an operator $\widehat{D} : R(\widehat{\Gamma}) \rightarrow R(\widehat{\Gamma})$,

$$\widehat{D}(\lambda_\gamma) = l(\gamma)\lambda_\gamma, \quad \gamma \in \Gamma.$$

We say that a quantum group \mathbb{G} acts on the dual of Γ by orientation preserving isometries if there exists a unital *-homomorphism $\alpha : C(\widehat{\Gamma}) \rightarrow C(\widehat{\Gamma}) \otimes C(\mathbb{G})$ such that

$$(\alpha \otimes \text{id}_{C(\mathbb{G})})\alpha = (\text{id}_{C(\widehat{\Gamma})} \otimes \Delta)\alpha$$

and moreover α restricts to a unital *-homomorphism $\alpha_0 : R(\widehat{\Gamma}) \rightarrow R(\widehat{\Gamma}) \odot R(\mathbb{G})$ satisfying the commutation relation

$$(3.1) \quad \alpha_0 \widehat{D} = (\widehat{D} \otimes \text{id}_{R(\mathbb{G})})\alpha_0$$

and preserving the canonical trace on $R(\widehat{\Gamma})$. For the motivation behind this definition and connections with spectral triples and noncommutative manifolds we refer to [BhS] and references therein.

Theorem 3.5 ([BhS]). *Let Γ be a discrete group with a fixed finite symmetric set of generators S . The category of all compact quantum groups acting on $\widehat{\Gamma}$ by orientation preserving isometries admits a universal (initial) object, denoted further by $QISO^+(\widehat{\Gamma}, S)$ and called the quantum group of orientation preserving isometries of $\widehat{\Gamma}$.*

When the choice of the generating set is clear, we write simply $QISO^+(\widehat{\Gamma})$. In particular if $\Gamma = \mathbb{F}_p$, the free group on p generators x_1, \dots, x_p , we use the generating set $S = \{x_1, x_1^{-1}, \dots, x_p, x_p^{-1}\}$. The following result is essentially a rephrasing of Theorem 5.1 of [BhS]; for the convenience of the reader we sketch the proof.

Theorem 3.6. *The quantum group $H^+(p, 0)$ is isomorphic to the quantum group of orientation preserving isometries of $\widehat{\mathbb{F}}_p$.*

Proof. Discussion after Theorem 2.6 of [BhS] implies that $\mathbb{G} := QISO^+(\mathbb{F}_p)$ is a compact matrix quantum group with a fundamental representation determined by a unitary $U = (U_{t,s})_{t,s \in S} \in M_{2p}(C(\mathbb{G}))$ such that the map

$$\alpha_0(\lambda_t) = \sum_{s \in S} \lambda_s \otimes U_{s,t}, \quad t \in S,$$

extends to a *-homomorphism $\alpha_0 : R(\widehat{\mathbb{F}}_p) \rightarrow R(\widehat{\mathbb{F}}_p) \odot R(\mathbb{G})$ satisfying (3.1). Relabel generators in S so that $S = \{x_{i\alpha}, i\alpha \in \mathcal{J}_p\}$ and $x_{i\alpha} = x_{i\alpha}^{-1}$ for each $i\alpha \in \mathcal{J}_p$. Then the fact that α_0 is a *-map implies that the entries of U satisfy the relations (2.2) (where we write $U_{i\alpha, j\beta} := U_{x_{i\alpha}, x_{j\beta}}$). Further unitality of α_0 (specifically the conditions $\alpha_0(x_t)\alpha_0(x_{t^{-1}}) = 1_{C(\widehat{\mathbb{F}}_p)} \otimes 1_{\mathbb{G}}$ for each $t \in S$) together with unitality of U imply that each $U_{i\alpha}$ is a partial isometry. Finally a combinatorial argument shows that no additional relations are implied by the fact that the *-homomorphism α_0 satisfies (3.1); hence the universal properties defining the standard fundamental representations of \mathbb{G} and $H^+(p, 0)$ coincide. \square

Finally we describe $A_h(p, q)$ in terms of the intertwiners between the tensor powers of the fundamental corepresentation.

Proposition 3.7. *The algebra $A_h(p, q)$ is the universal C^* -algebra generated by the entries of a unitary $2p + q$ by $2p + q$ matrix U such that the vector ξ defined in Proposition 2.2 is a fixed vector for $U^{\otimes 2}$ and the map $e_{i\alpha} \rightarrow e_{i\alpha} \otimes e_{i\alpha} \otimes e_{i\alpha}$, $e_M \rightarrow e_M \otimes e_M \otimes e_M$ defines a morphism in $\text{Hom}(U; U^{\otimes 3})$.*

Proof. The proof is very similar to that of Proposition 2.10. The difference lies in the fact that we obtain conditions of the type $(i\alpha, j\beta \in \mathcal{J}_p, M, N \in \{1, \dots, q\}, y, x \in \mathcal{J}_{p+q})$


$$\begin{aligned} U_{j\beta, i\alpha} U_{y, \bar{i}\alpha} U_{x, i\alpha} &= \delta_y^{\bar{j}\beta} \delta_x^{j\beta} U_{j\beta, i\alpha}, \\ U_{N, i\alpha} U_{y, \bar{i}\alpha} U_{x, i\alpha} &= \delta_y^N \delta_x^N U_{N, i\alpha}, \\ U_{j\beta, M} U_{y, M} U_{x, M} &= \delta_y^{\bar{j}\beta} \delta_x^{j\beta} U_{j\beta, M}, \\ U_{N, M} U_{y, M} U_{x, M} &= \delta_y^N \delta_x^N U_{N, M}. \end{aligned}$$

These together with relations (2.2)-(2.5) imply that all $U_{z,y}$ ($z, y \in \mathcal{J}_{p,q}$) are partial isometries. \square

4. CATEGORIES OF REPRESENTATIONS VIA PARTITIONS

In this section we describe the categories of representations of the quantum groups we are considering in this paper in terms of (marked) partitions. Let $P(k, l)$ ($k, l \in \mathbb{N}_0$) be the set of partitions between k upper points and l lower points. Given $\pi \in P(k, l)$ and two multi-indices $i = (i_1, \dots, i_k) \in \mathcal{J}_{p,q}^k$ and $j = (j_1, \dots, j_l) \in \mathcal{J}_{p,q}^l$, we define a number $\delta_\pi(\begin{smallmatrix} i \\ j \end{smallmatrix}) \in \{0, 1\}$ in the following way: first place the indices (i_1, \dots, i_k) and (j_1, \dots, j_l) respectively on the upper and lower points and then put $\delta_\pi(\begin{smallmatrix} i \\ j \end{smallmatrix}) = 1$ if in any block of π the upper and lower sequences of indices in $\mathcal{J}_{p,q}$, say $(x_1, \dots, x_r) \in \mathcal{J}_{p,q}^r$ and $(y_1, \dots, y_s) \in \mathcal{J}_{p,q}^s$, satisfy $x_1 = \bar{x}_2 = x_3 = \bar{x}_4 = \dots$,

$y_1 = \bar{y}_2 = y_3 = \bar{y}_4 = \dots$, $x_1 = y_1$, and $\delta_\pi(\dot{j}) = 0$ otherwise. Thus $\delta_\pi(\dot{j}) = 1$ if and only if indices in each block of the partition π have the following pattern:



Further consider the following operator in $B((\mathbb{C}^{2p+q})^{\otimes k}; (\mathbb{C}^{2p+q})^{\otimes l})$:

$$(4.1) \quad T_\pi(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1, \dots, j_l \in \mathcal{J}_{p,q}} \delta_\pi(\dot{j}) e_{j_1} \otimes \dots \otimes e_{j_l}, \quad (i_1, \dots, i_k) \in \mathcal{J}_{p,q}^k.$$

We denote by $P_2, P_{12}, P_{even} \subset P$ respectively the pairings, the singletons and pairings, and the partitions having even blocks. Let also $NC_x = NC \cap P_x$, for any $x \in \{., 2, 12, \text{even}\}$, where $NC \subset P$ denotes the set of all non-crossing partitions. It turns out that such collections of partitions, known to correspond to representations of the orthogonal, symmetric, bistochastic and hyperoctahedral (quantum) groups ([BaS]), can be also used to describe the categories in our two-parameter context. Indeed, one can check, similarly as it was done in [BaS], that for P_{12} and P_{even} the usual operations of tensoring, concatenation (with the appropriate multiplication factor added for deleted closed blocks), and turning the partition upside-down correspond to tensoring, composing and passing to the adjoint on the level of the associated operators T_π defined in (4.1). Note that such a statement fails when we consider the whole category P – this explains why the case of $S^+(p, q)$, to be discussed later on, cannot be included in the following theorem.

Theorem 4.1. *For $\mathbb{G} = O^+(p, q)$ (respectively, $\mathbb{G} = B^+(p, q), H^+(p, q)$) let U denote the fundamental representation introduced earlier. Then for all $k, l \in \mathbb{N}_0$*

$$\text{Hom}(U^{\otimes k}; U^{\otimes l}) = \text{span}(T_\pi | \pi \in D(k, l)),$$

with $D = NC_2$ (respectively, $D = NC_{12}, NC_{even}$).

Proof. In the one-parameter case ($p = 0$) the result was proved in [BaS]; it turns out that the methods used there can be easily adopted to our framework when we work with the quantum groups listed in the theorem. Details will be provided only for the case of $O^+(p, q)$, as the other two follow similarly.

In fact for $O^+(p, q)$ the result is a consequence of the general facts about $A_o(F)$. Indeed, let $F \in M_n(\mathbb{R})$ satisfy $F = F^t$ and $F^2 = 1$, and consider the algebra $A_o(F)$. Our claim is that we have $\text{Hom}(U^{\otimes k}; U^{\otimes l}) = \text{span}(T_\pi | \pi \in NC_2(k, l))$, where $T_\pi(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1, \dots, j_l} \delta_\pi(\dot{j}) e_{j_1} \otimes \dots \otimes e_{j_l}$, with $\delta_\pi(\dot{j}) \in \mathbb{R}$ being given by:

$$\delta_\pi \begin{pmatrix} i_1 & \dots & i_k \\ j_1 & \dots & j_l \end{pmatrix} = \prod_{(i_r i_s) \in \pi} F_{i_r i_s} \prod_{(j_r j_s) \in \pi} F_{j_r j_s} \prod_{(i_r j_s) \in \pi} \delta_{i_r j_s}$$

In other words, δ_π is obtained by taking the product of F_{ab} 's, over all “horizontal” strings of p , and of δ_{ab} 's, over all “vertical” strings of π (note that as we only consider here pair partitions, in fact only one delta factor will appear for any given block of the partition). Observe further that in the case of the matrix $F = F_{p,q}$ used for defining $O^+(p, q)$, we obtain the δ numbers in the statement; it therefore suffices to establish the claim.

The proof of the claim has two steps. First, we prove that $\pi \rightarrow T_\pi$ transforms the categorical operations for partitions into the categorical operations for the linear

maps. This is indeed clear for the tensor product and for the duality. Regarding now the composition axiom, the only problem might come from the closed circles that appear in the middle: but here we can use the formula $T_{\cup}T_{\cap} = (2p + q) \cdot \text{id}$, with $2p + q = \sum_{i,j \in \mathcal{J}_{p,q}} F_{ij}$.

Summarizing, the spaces $\text{span}(T_{\pi} | \pi \in NC_2(k, l))$ form a tensor category in the sense of Woronowicz. Now since this tensor category is generated by T_{\cap} (that is because NC_2 is generated by \cap) the corresponding Hopf algebra is the one obtained by using the relation $T_{\cap} \in \text{Fix}(U^{\otimes 2})$. And since $T_{\cap} = \sum F_{ij} e_i \otimes e_j$, this algebra is $A_o(F)$, and the proof is finished.

Note in passing, that when F is not necessarily real, the result remains the same, but with the δ numbers obtained by making the product of F_{ab} 's over "oriented" horizontal strings, and then the product of δ_{ab} 's over vertical strings.

The cases of $B^+(p, q)$ and $H^+(p, q)$ follow in a similar way, first adding the singletons and then considering all partitions with blocks of even size. \square

Remark 4.2. The corresponding result holds also for the classical versions of $O^+(p, q)$, $S^+(p, q)$ and $H^+(p, q)$ (see Section 6), with the non-crossing partitions replaced by all partitions; the proof follows in a similar way.

For $S^+(p, q)$ the above proof does not work: as already mentioned above, the problem is that, since some of the blocks have an odd number of entries, the composition axiom does not hold for the implementation provided by the operators T_{π} described in (4.1). We therefore need to provide a new combinatorial description of the category of representations. From Theorem 2.9 we know that $S^+(p, q) \approx H_p^+ \hat{\star} S_q^+$; hence we begin by considering separately the cases $p = 0$ and $q = 0$. When $p = 0$, the quantum group $S^+(0, q)$ is a quantum symmetric group with the usual fundamental representation, so the corresponding category is NC , with the standard implementation providing the isomorphism (there are no 'conjugate' coordinates to deal with), as shown in [Ban₁] or in [BaS]. The situation for $q = 0$ is more complicated, as the defining fundamental representation of $S^+(p, 0)$ corresponds rather to the 'sudoku' representation of H_p^+ than to the representation studied from the categorical point of view in [BBC₂] and [BaS]. We begin by defining the relevant category of 'bulleted' partitions.

Definition 4.3. Let (for $k, l \in \mathbb{N}_0$) $P_{\bullet}(k, l)$ denote the collection of all partitions between k upper and l lower points, such that each point is marked either with a black or a white circle and we identify the partitions which differ by 'mirrored markings' in some of the blocks, so that for example  equals  in $P_{\bullet}(2, 3)$. The category $P_{\bullet} = \bigcup_{k, l \in \mathbb{N}_0} P_{\bullet}(k, l)$ is defined similarly to P , with the result of the concatenation operation being non-zero only if the corresponding markings match, remembering that we are allowed to replace colours of markings in any block of any given partition.

Denote by NC_{\bullet} the subcategory of all bulleted non-crossing partitions.

Given $\pi \in P_{\bullet}(k, l)$ and multi-indices $i = (i_1, \dots, i_k) \in \mathcal{J}_p^k$, $j = (j_1, \dots, j_l) \in \mathcal{J}_p^l$ define a number $\delta_{\pi}(\begin{smallmatrix} i \\ j \end{smallmatrix}) \in \{0, 1\}$ in the following way: $\delta_{\pi}(\begin{smallmatrix} i \\ j \end{smallmatrix}) = 1$ if for any pair of vertices in a fixed block of the partition π the corresponding indices are equal if the bullets on the vertices have the same colour and are 'conjugate' if the colours of the respective bullets are different. Define further $\tilde{T}_{\pi} : (\mathbb{C}^{2p})^{\otimes k} \rightarrow (\mathbb{C}^{2p})^{\otimes l}$ by

the usual formula ($i_1, \dots, i_k \in \mathcal{J}_p$):

$$(4.2) \quad \tilde{T}_\pi(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1, \dots, j_l \in \mathcal{J}_p} \delta_\pi(i_j) e_{j_1} \otimes \dots \otimes e_{j_l}.$$

We are ready to formulate the result describing the representation theory of $S^+(p, 0)$.

Theorem 4.4. *Let U denote the fundamental representation of $S^+(p, 0)$ introduced in Definition 2.7. We have for all $k, l \in \mathbb{N}_0$*

$$\text{Hom}(U^{\otimes k} U^{\otimes l}) = \text{span}(\tilde{T}_\pi | \pi \in NC_\bullet(k, l)).$$

Proof. It is standard to check that NC_\bullet is a tensor category satisfying the properties needed to apply the Tannaka-Krein duality of [Wor₂] and that the map $\pi \rightarrow \tilde{T}_\pi$ transforms the natural operations (concatenation with deletion of the closed blocks, tensoring, turning upside-down) to the corresponding operations on the level of linear maps. Proposition 2.10 implies that the category of representations of $S^+(p, 0)$ is generated by the morphisms \tilde{T}_{π_1} and \tilde{T}_{π_2} associated with the partitions

$$\pi_1 = \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \circ \end{array}, \quad \pi_2 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \end{array}.$$

The partitions π_1 and π_2 generate the whole NC_\bullet . Indeed, it is well known that the analogous ‘non-bulleted’ partitions generate NC , and it suffices to notice that π_1 and π_2 can be first combined to obtain singletons and then the ‘exchange’ partition $\pi_3 = \begin{array}{c} \bullet \\ \circ \end{array}$ (the fact that \tilde{T}_{π_3} intertwines U is in a sense a consequence of the fact that entries of u are self-adjoint). Once we know that the category generated by π_1 and π_2 contains partitions of all shapes and we can exchange colours of markings at every vertex using π_3 the generation statement is clear.

The argument above shows that the category of representations of $S^+(p, 0)$ is at least as big as NC_\bullet ; so there is an inclusion $\mathbb{G} \supset S^+(p, 0)$, where \mathbb{G} is the compact quantum group arising as the Tannaka-Krein dual of NC_\bullet via the described implementation. Note that the inclusion $\mathbb{G} \supset S^+(p, 0)$ can be described as the surjection on the level of function algebras: $C(\mathbb{G}) \rightarrow C(S^+(p, 0))$. To conclude the proof, by using a standard Peter-Weyl argument, it suffices to show that for each $k \in \mathbb{N}$ the number $d_k := \text{card}(\text{Hom}(1; U^{\otimes k}))$ is equal to $\text{card}(NC_\bullet(0, k))$. The numbers d_k are expressed by the formula

$$d_k = h_{S^+(p, 0)}(\text{Tr}(U^{\otimes k})),$$

where $h_{S^+(p, 0)}$ denotes the Haar state. It follows from the proof of Theorem 2.9 that when we compute the trace of u we obtain the two copies of the elements which turn up as the trace of the top-left block of u in the ‘sudoku’ picture. By Theorem 7.2 and Corollary 7.4 of [BCS] we know that the corresponding single random variable has (with respect to the Haar state of H_p^+) the free Poisson distribution with parameter $\frac{1}{2}$. As the Haar state is obviously preserved by the isomorphism between $S^+(p, 0)$ and H_p^+ we have

$$(4.3) \quad d_k = 2^k \sum_{\pi \in NC(0, k)} \left(\frac{1}{2}\right)^{\nu(\pi)} = \sum_{\pi \in NC(0, k)} 2^{k-\nu(\pi)}$$

(where $\nu(\pi)$ denoted the number of blocks in the partition π and we used the formula for the moments of the free Poisson distribution with parameter t , see e.g. [NiS]). It remains to check that the number in (4.3) equals $\text{card}(NC_\bullet(0, k))$.

But this follows from the fact that for every non-bulleted block of a partition in $NC(0, k)$ which has l legs we have 2^{l-1} choices of the bullet pattern. \square

Remark 4.5. Again we have a corresponding statement for the classical version of $S^+(p, 0)$, with the category NC_\bullet replaced by P_\bullet .

Note that the categories of representations listed in Theorem 4.1 can also be described in terms of the bulleted partitions with the fixed colouring pattern consistent with the implementation in (4.1) (so for example if a block in a pair partition is ‘vertical’, both ends are given the same colour, and if it is ‘horizontal’, then we have a colour exchange).

‘Free product’ of categories described by partitions. Recall that for $p, q > 0$ we have $S^+(p, q) = S^+(p, 0) \hat{\star} S^+(0, q)$, and Theorem 4.4 together with the discussion before it describe the categories of representations of both $S^+(p, 0)$ and $S^+(0, q)$. Before we use this decomposition to provide a description of the category for $S^+(p, q)$, let us discuss a general free product framework.

If $\mathbb{G}_1, \mathbb{G}_2$ are compact quantum groups and $\mathbb{G} = \mathbb{G}_1 \hat{\star} \mathbb{G}_2$, although the knowledge of irreducible representations of \mathbb{G}_1 and \mathbb{G}_2 suffices to describe the irreducible representations of \mathbb{G} , the description on the level of categories of representations seems to be quite difficult. If however respective categories for \mathbb{G}_1 and \mathbb{G}_2 , denoted say by \mathcal{C}_1 and \mathcal{C}_2 , are given by (non-crossing) partitions, there is a natural candidate for the category corresponding to \mathbb{G} – it should be given by all partitions which decompose into blocks of two colours, where the sub-partition obtained by looking only at the blocks of the first colour belongs to \mathcal{C}_1 and that for the second colour belongs to \mathcal{C}_2 . The tensoring and reflection operations are defined as for usual partitions, and so is the concatenation, with the caveat that to obtain a non-zero outcome the colours must fit together. The resulting category will be denoted by $\mathcal{C}_1 \star \mathcal{C}_2$ (note that the construction works well also for bulleted partitions). As in the proof of Theorem 4.4, using the Tannakian duality we can establish the existence of a compact quantum group \mathbb{G}' , whose category of representations equals $\mathcal{C}_1 \star \mathcal{C}_2$, and deduce the existence of the inclusion $\mathbb{G}' \supset \mathbb{G}$. To show that this inclusion is in fact an equality, we need to compare the dimensions of the fixed point spaces (or, equivalently, the moments of the laws describing the characters of the defining fundamental representations). As the Haar state of \mathbb{G} was shown in [Wan₁] to be the free product of the Haar states of \mathbb{G}_1 and \mathbb{G}_2 , the law related to \mathbb{G} arises as the free convolution of the corresponding distributions for \mathbb{G}_1 and \mathbb{G}_2 . Thus if this free convolution can be easily computed, and the categories \mathcal{C}_1 and \mathcal{C}_2 are identical, the corresponding count can be performed without much difficulty. In particular using the natural ‘free product’ implementations $\pi \rightarrow \hat{T}_\pi$ defined in the spirit of (4.1) we obtain the following result.

Theorem 4.6. *Let $n, m \in \mathbb{N}$. The category corresponding to the standard fundamental representations of $S_n^+ \hat{\star} S_m^+$ (respectively, $H_n^+ \hat{\star} H_m^+$, $B_n^+ \hat{\star} B_m^+$ and $O_n^+ \hat{\star} O_m^+$) is equal to the span of $\{\hat{T}_\pi : \pi \in \mathcal{C} \star \mathcal{C}\}$, where $\mathcal{C} = NC$ (respectively, NC_{12} , NC_{even} and NC_2).*

Proof. The proof follows as in Theorem 4.4, so we only provide the argument for the equality of the cardinality of partitions in $(\mathcal{C} \star \mathcal{C})(0, k)$ and the dimension of the fixed point spaces for tensor powers of the defining fundamental representations. Fix $x \in \{., 2, 12, \text{even}\}$ and for each $t > 0$ consider the probability measure μ_t

having as moments $\int \lambda^k d\mu_t(\lambda) = \sum_{\pi \in NC_x} t^{\nu(\pi)}$, where $\nu(\pi)$ denotes the number of blocks in π . The results in [BaS] imply that μ_1 is the measure describing the distribution of the character of the defining representation of the quantum group corresponding to x , and also that the measures $\{\mu_t : t > 0\}$ form a semigroup with respect to the free convolution \boxplus (see [Voi]). A standard argument using the definition of the free product construction implies that the character of the defining representation for $\mathbb{G}_n^+ \hat{\star} \mathbb{G}_m^+$ follows the law $\mu_1 \boxplus \mu_1 = \mu_2$, with moments given by $\int \lambda^k d\mu_2(\lambda) = \sum_{\pi \in NC_x} 2^{\nu(\pi)}$. But this is exactly the number of partitions featuring in $(\mathcal{C} \star \mathcal{C})(0, k)$ – using the surjective map $\mathcal{C} \star \mathcal{C} \rightarrow \mathcal{C}$ we see that each element in the preimage of a given $\pi \in \mathcal{C}(0, k)$ is the partition π with each block given one of the two colours; thus the preimage has $2^{\nu(\pi)}$ elements. \square

Before we prove an analogous, more technical, result for the quantum group $S^+(p, q)$ isomorphic to $S^+(p, 0) \hat{\star} S^+(0, q)$ we need a simple observation related to the free cumulants of probability measures. All relevant definitions can be found in Lectures 11 and 12 in [NiS].

Lemma 4.7. *Assume that ν is a compactly supported probability measure on \mathbb{R} . Let X be a real-valued random variable with distribution ν , let $\mu = \nu \boxplus \nu$ and let ν' be the law of $2X$. Denote the respective free cumulants of μ and of ν' by $(\kappa_k(\mu))_{k=1}^\infty$ and $(\kappa_k(\nu'))_{k=1}^\infty$. Then for each $k \in \mathbb{N}$*

$$\kappa_k(\nu') = 2^{k-1} \kappa_k(\mu).$$

Proof. It follows from Proposition 12.3 in [NiS] that $\kappa_n(\nu) = \frac{1}{2} \kappa_n(\mu)$. Further the fact that the moments and cumulants are related by the Speicher's moment-cumulant formula ([Spe]):

$$(4.4) \quad d_k = \sum_{\pi \in NC(0, k)} \prod_{b: \text{block in } \pi} \kappa_{\text{card}(b)},$$

implies that $\kappa_k(\nu') = 2^k \kappa_k(\nu)$. \square

The following theorem completes the description of the categories for the main family of quantum groups studied in this paper.

Theorem 4.8. *Let u denote the fundamental representation of $S^+(p, q)$ introduced in Definition 2.7. Then for all $k, l \in \mathbb{N}_0$*

$$\text{Hom}(U^{\otimes k}; U^{\otimes l}) = \text{span}(\tilde{T}_\pi | \pi \in (NC_\bullet \star NC)(k, l)).$$

Proof. As in the proof of Theorem 4.6 we only provide the argument for the equality of the number of partitions in $(NC_\bullet \star NC)(0, k)$ and the k -th moment of the character of the defining representation of $S^+(p, q)$. The latter can be expressed via the moment-cumulant formula (4.4), with the free cumulants $(\kappa_k)_{k=1}^\infty$ given by the sum of the free cumulants of the corresponding laws of the characters for $S^+(p, 0)$ and $S^+(0, q)$ (this is a consequence of the discussion before Theorem 4.6 and Proposition 12.3 in [NiS]). The free cumulants for the second law (which is the free Poisson distribution) are equal to 1. As the free Poisson laws form a free convolution semigroup, the proof of Theorem 4.4 and Lemma 4.7 imply that the free cumulants for the first law are equal to 2^{k-1} . Hence the moment-cumulant formula for the k -th moment of the law we are interested in yields:

$$d_k = \sum_{\pi \in NC(0, k)} \prod_{b: \text{block in } \pi} (2^{\text{card}(b)-1} + 1).$$

It remains to note that the above number indeed corresponds to the number of partitions in $(NC_\bullet \star NC)(0, k)$: for each block b in a given non-crossing partition π of k -points we can choose whether it ‘comes’ from NC_\bullet or NC and in the first case we have additionally $2^{\text{card}(b)-1}$ choices of inequivalent bulleting patterns. \square

5. RELATIONS BETWEEN THE TWO-PARAMETER FAMILIES

Recall the diagram (1.2) describing the relations between O_n^+ , B_n^+ , H_n^+ and S_n^+ . Universal properties imply that the corresponding diagram can be drawn in the two-parameter case; for each $p, q \in \mathbb{N}_0$ we have

$$(5.1) \quad \begin{array}{ccc} A_o(p, q) & \rightarrow & A_b(p, q) & & O^+(p, q) & \supset & B^+(p, q) \\ & & \downarrow & & \cup & & \cup \\ & & A_h(p, q) & \rightarrow & A_s(p, q) & & H^+(p, q) \supset S^+(p, q) \end{array}$$

Below we describe further connections between the quantum groups studied above, first discussing the general framework.

Definition 5.1. Assume that we have surjective morphisms of Hopf C^* -algebras, corresponding to inclusions of compact quantum groups, as follows:

$$(5.2) \quad \begin{array}{ccccc} A & \rightarrow & B & & \mathbb{G} & \supset & \mathbb{H} \\ & & \downarrow & & \downarrow & & \cup & & \cup \\ & & C & \rightarrow & D & & \mathbb{K} & \supset & \mathbb{L} \end{array}$$

- (1) We write $\mathbb{G} = \langle \mathbb{H}, \mathbb{K} \rangle$ if the $*$ -algebra ideal $\ker(A \rightarrow B) \cap \ker(A \rightarrow C)$ contains no nonzero Hopf ideal.
- (2) We write $\mathbb{L} = \mathbb{H} \cap \mathbb{K}$ when $\ker(A \rightarrow D)$ is the Hopf ideal generated by $\ker(A \rightarrow B)$ and $\ker(A \rightarrow C)$.

The definitions above can be easily seen to coincide with the standard ones in the classical case.

More generally, given morphisms $\alpha : A \rightarrow B$ and $\beta : A \rightarrow C$ as above, one can define a quantum group $\langle \mathbb{G}, \mathbb{H} \rangle \subset \mathbb{K}$ by the formula $C(\langle \mathbb{G}, \mathbb{H} \rangle) = A/I$, where I is the biggest Hopf ideal contained in $\ker(\alpha) \cap \ker(\beta)$. Once again, this definition agrees with the usual one in the classical case.

These notions are best understood in terms of the Hopf image formalism, introduced in [BaB]. Consider indeed the morphism $(\alpha, \beta) : C \rightarrow A \times B$. Then $C(\langle \mathbb{G}, \mathbb{H} \rangle)$ is the Hopf image of (α, β) , and we have $\langle \mathbb{G}, \mathbb{H} \rangle = \mathbb{K}$ if and only if (α, β) is inner faithful.

Similarly given a 4-term diagram as in the above definition, we can form the Hopf ideal $J = \langle \ker(A \rightarrow B), \ker(A \rightarrow C) \rangle$, and define a quantum group $\mathbb{H} \cap \mathbb{K}$ by $C(\mathbb{H} \cap \mathbb{K}) = A/J$. We have $\mathbb{L} = \mathbb{H} \cap \mathbb{K}$ if and only if the above condition (2) is satisfied.

In order to deal effectively with part (1) of Definition 5.1, we use the following Tannakian reformulation.

Lemma 5.2. *Consider the diagram (5.2) and denote the Hopf morphism between A and B (respectively, between A and C) by α (respectively, β). We have $\mathbb{K} = \langle \mathbb{G}, \mathbb{H} \rangle$ if and only if*

$$\text{Fix}(R) = \text{Fix}((\text{id} \otimes \alpha)R) \cap \text{Fix}((\text{id} \otimes \beta)R)$$

for any representation R arising as a tensor product between U and \bar{U} 's, with U being the fundamental representation of \mathbb{G} .

Proof. The result is a consequence of general considerations in [BaB]; we sketch the proof below.

Let I be the biggest Hopf ideal contained in $\ker(\alpha) \cap \ker(\beta)$, whose existence follows from [BaB], and let $\mathbb{K}_1 = \langle \mathbb{G}, \mathbb{H} \rangle$ be the compact quantum group given by $C(\mathbb{K}_1) = C(\mathbb{K})/I$.

By Frobenius duality, the collection of equalities in the statement is equivalent to the following collection of equalities:

$$\text{Hom}(R; S) = \text{Hom}((\text{id} \otimes \alpha)R; (\text{id} \otimes \alpha)S) \cap \text{Hom}((\text{id} \otimes \beta)R; (\text{id} \otimes \beta)S),$$

where now both R and S are representations arising as tensor products of several copies of U and \bar{U} .

According to the general results of Woronowicz in [Wor₂], the Hom spaces on the right form a Tannakian category, which should therefore correspond to a certain compact quantum group \mathbb{K}_2 . Our claim is that we have $\mathbb{K}_1 = \mathbb{K}_2$. Indeed, this follows from Theorem 8.4 of [BaB] applied to the morphism $(\alpha, \beta) : C \rightarrow A \times B$ (and can be viewed as a consequence of the general Peter-Weyl type results in [Wor₁] and the Tannakian duality results in [Wor₂]).

In particular $\mathbb{K} = \mathbb{K}_1$ is equivalent to $\mathbb{K} = \mathbb{K}_2$, and we are done. \square

Before we formulate the specific intersection/generation results in the cases in which we are interested, we need another lemma:

Lemma 5.3. *Let \mathbf{A} be a C^* -algebra, $n \in \mathbb{N}$ and let $U \in M_n(\mathbf{A})$ be a unitary matrix whose entries are partial isometries. If for each $z \in \{1, \dots, n\}$ there is $\sum_{y=1}^n U_{y,z} = 1$, then each $U_{y,z}$ is an orthogonal projection.*

Proof. Fix $y, z \in \{1, \dots, n\}$. We have $U_{y,z} = 1 - \sum_{x \neq y} U_{x,z}$. Proposition 3.2 implies that multiplying the last equality by $U_{y,z}^*$ on the right yields $U_{y,z}U_{y,z}^* = U_{y,z}^*$. Hence $U_{y,z}$ is a selfadjoint projection. \square

Theorem 5.4. *The following relations hold:*

- (1) $O^+(p, q) = \langle B^+(p, q), H^+(p, q) \rangle$.
- (2) $S^+(p, q) = B^+(p, q) \cap H^+(p, q)$.

Proof. (1) follows from Lemma 5.2, Theorem 4.1, and the equality $NC_2 = NC_{12} \cap NC_{\text{even}}$.

(2) is a direct consequence of Lemma 5.3. \square

Free products. We recall from Section 2 that we have a canonical isomorphism $A_o(p, q) \simeq A_o(2p+q)$. It is well known that for any $n, m \in \mathbb{N}$ there exists a natural surjective map $A_o(m+n) \rightarrow A_o(m) \star A_o(n)$. Similar maps turn out to exist in all the cases considered in this paper.

Proposition 5.5. *Let $g \in \{o, b, h, s\}$. There exists a natural surjective map $A_g(p, q) \rightarrow A_g(p, 0) \star A_g(0, q)$ such that the following diagram (in which the vertical maps are the ones introduced earlier in this section) is commutative:*

$$\begin{array}{ccc} A_o(p, q) & \rightarrow & A_o(p, 0) \star A_o(0, q) \\ \downarrow & & \downarrow \\ A_g(p, q) & \rightarrow & A_g(p, 0) \star A_g(0, q) \end{array}$$

The horizontal maps in the diagram above intertwine respective comultiplications.

Proof. Let $U_1 \in M_{2p}(A_g(p, 0))$ and $U_2 \in M_q(A_g(0, q))$ be fundamental unitary representations of the respective compact quantum groups. Consider the unitary matrix $\begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix}$ viewed as an element of $M_{2p+q}(A_g(p, 0) \star A_g(0, q))$. It is easy to see that it satisfies all the conditions required of the generating matrix in $M_{2p+q}(A_g(p, q))$, so the universality implies the existence of a *-homomorphism requested by the proposition. It is easily seen to be surjective, as entries of U_1 (respectively, U_2) generate $A_g(p, 0)$ (respectively, $A_g(0, q)$). An explicit description of the comultiplications involved implies the last statement of the proposition.

Recall that the existence of surjective (compact quantum group) morphisms from $A_o(p, q)$ to $A_g(p, q)$ followed in an analogous way; the fact that the diagram above is commutative is thus a direct consequence of the definitions of the maps considered. \square

Using directly the language of compact quantum groups we obtain the following corollary:

Corollary 5.6. *Let $\mathbb{G} \in \{O^+, H^+, B^+, S^+\}$. There exist natural inclusions*

$$\begin{array}{ccc} O^+(p, q) & \supset & O^+(p, 0) \hat{\star} O^+(0, q) \\ \cup & & \cup \\ \mathbb{G}(p, q) & \supset & \mathbb{G}(p, 0) \hat{\star} \mathbb{G}(0, q) \end{array}$$

In the case of $A_o(p, q)$ the map described in Proposition 5.5 and the isomorphisms/surjections recalled before it can be combined into the following commutative diagram:

$$\begin{array}{ccc} A_o(p, q) & \rightarrow & A_o(p, 0) \star A_o(0, q) \\ \downarrow & & \downarrow \\ A_o(2p+q) & \rightarrow & A_o(2p) \star A_o(q) \end{array}$$

(the commutativity is the consequence of the fact how the isomorphism was described in the proof of Theorem 2.1 and the block-diagonal form of the matrices $C_{p,q}$ implementing it).

Theorem 2.9 implies that the inclusion $S^+(p, q) \supset S^+(p, 0) \hat{\star} S^+(0, q)$ is in fact an isomorphism. In all the other cases when both p and q are non-zero (for $B^+(p, q)$ we actually need to assume $q > 1$ to avoid trivialities) $\mathbb{G}(p, 0) \hat{\star} \mathbb{G}(0, q)$ is a proper quantum subgroup of $\mathbb{G}(p, q)$. This is the content of the next proposition.

Proposition 5.7. *If $p, q \neq 0$ then the inclusions $O^+(p, 0) \hat{\star} O^+(0, q) \subset O^+(p, q)$ and $H^+(p, 0) \hat{\star} H^+(0, q) \subset H^+(p, q)$ are proper. If $p > 0$ and $q > 1$ then the inclusion $B^+(p, 0) \hat{\star} B^+(0, q) \subset B^+(p, q)$ is proper.*

Proof. It is enough to show that the corresponding homomorphisms $A_o(p, q) \rightarrow A_o(p, 0) \star A_o(0, q)$, $A_h(p, q) \rightarrow A_h(p, 0) \star A_h(0, q)$ and $A_b(p, q) \rightarrow A_b(p, 0) \star A_b(0, q)$ are not injective; in other words it suffices to find unitary matrices satisfying the defining conditions for $A_o(p, q)$, $A_h(p, q)$ and $A_b(p, q)$ which are not of the form $\begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix}$ with U_1 a $2p$ by $2p$ matrix. In the case of the orthogonal and the bistochastic group the existence of such matrices is visible already at the commutative level.

Consider then the case of H^+ . It suffices to find suitable matrices for $p = q = 1$. Let then Q_A, Q_B, P_A, P_B be non-zero orthogonal projections on some Hilbert space \mathbb{H} such that $Q_A + Q_B + P_A + P_B = 1$ and let A, B, C, D be partial isometries in $B(\mathbb{H})$ such that $AA^* = Q_A$, $A^*A = P_A$, $BB^* = Q_B$, $B^*B = P_B$, $CC^* = 1 - C^*C = P_B + Q_A$, $DD^* = 1 - D^*D = P_A + Q_B$. The existence of such partial isometries can be assured by choosing all the projections to have infinite-dimensional ranges. Then the matrix

$$\begin{bmatrix} A & B & C \\ B^* & A^* & C^* \\ D & D^* & 0 \end{bmatrix}$$

can be checked to be a unitary satisfying assumptions listed in Definition 3.1. \square

Note that the above construction excludes the possibility of the partial isometries A, B, C, D generating a commutative algebra. This is a general fact – in Theorem 6.1 below we will see that if we additionally request commutativity then the ‘off-diagonal’ terms of a unitary satisfying the defining conditions of $A_h(p, q)$ must vanish.

Quantum permutation groups as quotient quantum groups of $H^+(p, q)$.

Definition 5.8. If \mathbb{G}_1 and \mathbb{G}_2 are compact quantum groups and $\pi : C(\mathbb{G}_1) \rightarrow C(\mathbb{G}_2)$ is a unital injective $*$ -homomorphism intertwining the respective coproducts, then \mathbb{G}_1 is said to be a quotient quantum group of \mathbb{G}_2 (alternatively, \mathbb{G}_2 is said to be a quantum group extension of \mathbb{G}_1).

Consider once again the quantum group $H^+(p, q)$. As the generators of $A_h(p, q)$, denoted by $U_{z,y}$ ($z, y \in \mathcal{J}_{p,q}$) are partial isometries, the operators $P_{z,y} := U_{z,y}^* U_{z,y}$ are projections.

Theorem 5.9. *The C^* -subalgebra of $A_h(p, q)$ generated by $P_{z,y}$ ($z, y \in \mathcal{J}_{p,q}$) is isomorphic to $A_s(2p + q)$.*

Proof. Let A denote the C^* -subalgebra of $A_h(p, q)$ generated by $P_{z,y}$ ($z, y \in \mathcal{J}_{p,q}$). Note first that the generators $\{P_{z,y} : z, y \in \mathcal{J}_{p,q}\}$ satisfy the same relations as generators of $A_s(2p + q)$; indeed for $z, y \in \mathcal{J}_{p,q}$

$$\begin{aligned} \sum_{x \in \mathcal{J}_{p,q}} P_{z,x} P_{y,x} &= \sum_{x \in \mathcal{J}_{p,q}} U_{z,x}^* U_{z,x} U_{y,x}^* U_{y,x} = \delta_z^y \sum_{x \in \mathcal{J}_{p,q}} U_{z,x}^* U_{z,x} U_{z,x}^* U_{z,x} \\ &= \delta_z^y \sum_{x \in \mathcal{J}_{p,q}} U_{z,x}^* U_{z,x} = \delta_z^y \sum_{\bar{x} \in \mathcal{J}_{p,q}} U_{\bar{z}, \bar{x}} U_{\bar{z}, \bar{x}}^* = \delta_z^y 1. \end{aligned}$$

Similarly it follows from Corollary 3.3 that

$$\begin{aligned} \sum_{x \in \mathcal{J}_{p,q}} P_{x,z} P_{x,y} &= \sum_{x \in \mathcal{J}_{p,q}} U_{x,z}^* U_{x,z} U_{x,y}^* U_{x,y} = \delta_z^y \sum_{x \in \mathcal{J}_{p,q}} U_{x,z}^* U_{x,z} U_{x,z}^* U_{x,z} \\ &= \delta_z^y \sum_{x \in \mathcal{J}_{p,q}} U_{x,z}^* U_{x,z} = \delta_z^y 1. \end{aligned}$$

If Δ denotes the coproduct of $A_h(p, q)$ then for $z, y \in \mathcal{J}_{p,q}$

$$\Delta(P_{z,y}) = \sum_{x \in \mathcal{J}_{p,q}} P_{z,x} \otimes P_{x,y}.$$

Hence A is isomorphic to $C(\mathbb{G})$, where \mathbb{G} is a quantum subgroup of S_{2p+q}^+ . It remains to check that \mathbb{G} is actually S_{2p+q}^+ . For that it suffices to analyse the corresponding categories of representations, and even more specifically to check that cardinalities of fixed point sets of tensor powers of the fundamental representations of \mathbb{G} coincide with those appearing for S_{2p+q}^+ (see Section 4, where this method was applied several times).

In the partition picture described in Section 4 the passage from $H^+(p, q)$ to \mathbb{G} corresponds to considering only partitions with even number of upper and lower points and composing them (both from above and from below) with a suitable number of copies of the partition \updownarrow . This means that we are left only with these partitions in $NC(2k, 2l)$ in which the pairs of points beginning at an odd place are necessarily joined (it is easy to see that all such partitions arise in this procedure); we denote them by $NC_{\text{join}}(2k, 2l)$. Now collapsing the pairs listed above we obtain a bijection between $NC_{\text{join}}(2k, 2l)$ and $NC(k, l)$. This suffices to perform the count needed to finish the proof. \square

Using the dual language and noticing that the map $A_p(p, q) \rightarrow A$ is an injective morphism preserving the respective coproducts, we obtain the following corollary.

Corollary 5.10. *The quantum group $H^+(p, q)$ is an extension of S_{2p+q}^+ .*

6. CLASSICAL VERSIONS AND INTERPRETATIONS IN TERMS OF CLASSICAL/QUANTUM SYMMETRIES

Each of the quantum groups studied above has a classical version. These are understood as follows: if \mathbb{G} is a compact quantum group then the quotient of the algebra $C(\mathbb{G})$ by its commutator ideal is isomorphic to the algebra of continuous functions on a certain uniquely determined compact group G . Then we call G the classical version of \mathbb{G} . Note that if we use the notation $G(p, q)$ for the classical version of the quantum group $G^+(p, q)$, then Corollary 5.6 and a straightforward analysis of classical versions of free products described in Section 1 implies that we have the following inclusions:

$$\begin{array}{ccc} O(p, q) & \supset & O(p, 0) \times O(0, q) \\ \cup & & \cup \\ G(p, q) & \supset & G(p, 0) \times G(0, q) \end{array}$$

The combination of results from the previous section and [BaS] allow us to identify the classical version of $O^+(p, q)$ with O_{2p+q} , the classical version of $B^+(p, q)$

with $O_{2p+q-i(p,q)}$ (where $i(p,q) = 1$ if $pq = 0$ and $i(p,q) = 2$ if $p, q > 0$) and the classical version of $S^+(p,q)$ with $H_p \times S_q$.

Theorem 6.1. *The classical version of $H^+(p,q)$ is $(\mathbb{T}^p \rtimes H_p) \times H_q$ (recall that $\mathbb{T}^p \rtimes H_p$ is the usual isometry group of \mathbb{T}^p).*

Proof. Observe first that if $\{U_{z,y} : z, y \in \mathcal{J}_{p,q}\}$ are commuting partial isometries which satisfy the condition (2.6) and form a unitary matrix, then $U_{i\alpha, M} = U_{M, i\alpha} = 0$ for all $i\alpha \in \mathcal{J}_p$ and $M \in \{1, \dots, q\}$. Indeed, the projections $U_{i\alpha, M}^* U_{i\alpha, M}$ and $U_{i\alpha, M}^* U_{i\alpha, M} = U_{i\alpha, M} U_{i\alpha, M}^*$ are then equal; as due to Proposition 3.2 they are pairwise orthogonal to each other, they must vanish (similar argument applies to $U_{M, i\alpha}$). Hence to prove the theorem it suffices to consider separately cases $p = 0$ and $q = 0$. The first one was treated in [BBC₂].

Assume then that $q = 0$. The result can be deduced from calculations in [Bho], but we can also offer an explicit isomorphism. Denote the images of elements of $\{U_{i\alpha, j\beta} : i\alpha, j\beta \in \mathcal{J}_p\}$ in the quotient space (with respect to the commutator ideal of $A_h(p, 0)$ by $\{\hat{U}_{i\alpha, j\beta} : i\alpha, j\beta \in \mathcal{J}_p\}$ and let the quotient C^* -algebra be denoted by $A_h^{\text{com}}(p, 0)$. Identify $C(\mathbb{T}^p \rtimes H_p)$ as a C^* -algebra in a natural way with $C(\mathbb{T}^p) \otimes C(S_p) \otimes C(\underbrace{\mathbb{Z}_2 \times \dots \times \mathbb{Z}_2}_{p \text{ times}})$. Let $\chi_{\alpha, \beta} \in C(S_p)$ be the characteristic

function of the set of these permutations which map α into β and let $\kappa_{0, \alpha}, \kappa_{1, \alpha} \in C(\mathbb{Z}_2 \times \dots \times \mathbb{Z}_2)$ be characteristic functions of sets which have respectively 0 or 1 in the α -coordinate. One can then check that the map

$$\hat{U}_{0\alpha, 0\beta} \rightarrow z_\alpha \otimes \chi_{\alpha, \beta} \otimes \kappa_{0, \alpha},$$

and

$$\hat{U}_{0\alpha, 1\beta} \rightarrow z_\alpha \otimes \chi_{\alpha, \beta} \otimes \kappa_{1, \alpha}$$

extends to an isomorphism of $A_h^{\text{com}}(p, 0)$ with $C(\mathbb{T}^p \rtimes H_p)$; this isomorphism preserves also the respective coproducts. \square

Note that from the point of view of the philosophy presented in [BaS] $H^+(p, 0)$ can therefore be viewed as the liberation of a group of isometries of p -copies of the circle. This is consistent with the results of Section 5 of [BhS], where $H^+(p, 0)$ was first discovered as the quantum isometry group of $C(\widehat{\mathbb{F}}_p)$ - in other words, of the ‘free torus’ (note that $C(\widehat{\mathbb{F}}_p)$ is the universal C^* -algebra generated by p unitaries, whereas $C(\mathbb{T}^p)$ is the universal C^* -algebra generated by p commuting unitaries). Similarly the quantum group O_p^+ can be viewed as the quantum symmetry group of \mathbb{S}^{2p} , the free sphere studied in [BaG]. To strengthen this analogy we observe the following fact.

Proposition 6.2. *There is a natural surjection from $C(\mathbb{S}^{2p})$ to $C(\widehat{\mathbb{F}}_p)$.*

Proof. Recall that $C(\mathbb{S}^{2p})$ is the universal C^* -algebra generated by selfadjoint operators $\{x_i, y_i : i = 1, \dots, p\}$ such that $\sum_{i=1}^p x_i^2 + y_i^2 = 1$ and $C^*(\mathbb{F}_p) \approx C(\widehat{\mathbb{F}}_p)$ is the universal C^* -algebra generated by unitaries $\{u_i : i = 1, \dots, p\}$. It suffices to observe that the quotient of $C(\mathbb{S}^{2p})$ by the closed two-sided ideal generated by expressions $x_i y_i = y_i x_i, x_i^2 + y_i^2 = \frac{1}{p}$ ($i = 1, \dots, p$) is naturally isomorphic to $C(\widehat{\mathbb{F}}_p)$ - the isomorphism maps images of x_i and y_i in the quotient respectively to $\frac{1}{2\sqrt{p}}(u_i + u_i^*)$ and $\frac{i}{2\sqrt{p}}(u_i - u_i^*)$. \square

7. QUANTUM GROUP $H_s^+(p, q)$

It is natural to ask whether the two-parameter families studied in this paper in a sense exhaust all possible natural choices in the ‘ $F_{p,q}$ ’ framework. When we studied $A_h(p, q)$ and $A_s(p, q)$ we required the entries of the fundamental corepresentation U to satisfy respectively the conditions of the type $x = xx^*x$ and $x = x^*x$, which have natural descriptions in terms of the intertwiners between U and its tensor powers (see Proposition 2.10 and 3.7). We could equally well describe in terms of the intertwiners conditions on the entries of the type say $x = xx^*xx^*$. These introduce only one new possibility, as explained by the next proposition.

Proposition 7.1. *Let x be a bounded operator on a Hilbert space, $k \in \mathbb{N}$. If $x = (x^*x)^k$ or $x = (xx^*)^k$ then x is an orthogonal projection. If $x = x(x^*x)^k$, then x is a partial isometry. If $x = x^*(xx^*)^k$, then x is a partial symmetry (a selfadjoint partial isometry).*

Proof. We only prove the last statement, leaving the first two to the reader. Suppose $x = x^*(xx^*)^k$. Then $x^* = x(x^*x)^k$, so $x^2 = (xx^*)^{k+1}$ and $(x^*)^2 = (x^*x)^{k+1}$. Taking the adjoint of the last relation we obtain that $(x^*x)^{k+1} = (xx^*)^{k+1}$. As both xx^* and x^*x are positive, the last equality holds if and only if x is normal. Now it remains to observe that for $z \in \mathbb{C}$ the equality $z = \bar{z}|z|^k$ holds if and only if $z \in \{-1, 0, 1\}$ and apply the spectral theorem to end the proof. \square

The above proposition motivates the next definition.

Definition 7.2. The algebra $A_{hs}(p, q)$ is the universal C^* -algebra generated by partial symmetries $\{U_{z,y} : z, y \in \mathcal{J}_{p+q}\}$ which satisfy all the relations required of generators of $A_o(p, q)$.

Proposition 7.3. *The algebra $A_{hs}(p, q)$ is the universal C^* -algebra generated by two families of partial symmetries $\{U_{i\alpha, j\beta} : i\alpha, j\beta \in \mathcal{J}_p\}$ and $\{U_{M,N} : M, N \in \{1, \dots, q\}\}$, such that both matrices $(U_{i\alpha, j\beta})_{i\alpha, j\beta \in \mathcal{J}_p}$ and $(U_{M,N})_{M, N=1}^q$ are magic unitaries and moreover*

$$(7.1) \quad U_{i\alpha, j\beta} = U_{i\alpha, \bar{j}\beta}, \quad i\alpha, j\beta \in \mathcal{J}_p.$$

Proof. It suffices to show that whenever $i\alpha \in \mathcal{J}_p$ and $M \in \{1, \dots, q\}$ then $U_{i\alpha, M} = 0 = U_{M, i\alpha}$. Due to Proposition 3.2 we have $(U_{i\alpha, M})^2 = 0 = (U_{M, i\alpha})^2 = 0$, and the proof is finished. \square

Before we formulate the main result we need to recall the definition of a ‘higher-order’ quantum hyperoctahedral group first defined in [BBC₂] and later studied for example in [BCS] and [BBCC].

Definition 7.4. Denote by $A_h^{(4)}(n)$ the universal C^* -algebra generated by the entries of an n by n unitary U , such that \bar{U} is also unitary, for each $i, j = 1, \dots, n$ the entry U_{ij} is normal, and $U_{ij}^4 = U_{ij}U_{ij}^*$ is an orthogonal projection. When $U \in M_n \otimes A_s(n)$ is interpreted as the fundamental unitary corepresentation, we view $A_h^{(4)}(n)$ as the algebra of continuous functions on the quantum hyperoctahedral group of order 4 on n coordinates, H_n^{4+} .

Note that if in the above definition 4 is replaced by 2 we obtain the quantum hyperoctahedral group H_n^+ introduced in Definition 1.4. The factors 2 and 4 can be replaced by arbitrary $s \in \mathbb{N}$, as shown in the papers cited above.

Theorem 7.5. *The algebra $A_{hs}(p, q)$ is the algebra of continuous functions on a compact quantum group, denoted further $H_s^+(p, q)$. The unitary $U = (U_{z,y})_{z,y \in \mathcal{J}_{p,q}} \in M_{2p+q} \otimes A_{hs}(p, q)$ is the fundamental representation of $S^+(p, q)$. The algebra $A_{hs}(p, q)$ is isomorphic to the free product $A_{hs}(p, 0) \star A_h(q)$; on the level of quantum groups we have $H_s^+(p, q) \approx H_p^{4+} \star H_q^+$.*

Proof. By Proposition 7.3 it suffices to consider separately the cases $p = 0$ and $q = 0$. The fact that $A_{hs}(0, q) \approx A_h(q)$ (with the isomorphism preserving natural fundamental corepresentations) is immediate. Assume then that $q = 0$. Note that the fundamental corepresentation has then the form

$$(7.2) \quad \begin{bmatrix} p_A - q_A & p_B - q_B & p_C - q_C & p_D - q_D & \cdots \\ p_B - q_B & p_A - q_A & p_D - q_D & p_C - q_C & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

where for each $X \in \{A, B, C, D, \dots\}$ the pair (p_X, q_X) is a pair of mutually orthogonal projections.

Following the permutation procedure described in the proof of Theorem 2.9 we can describe the fundamental representation as a ‘sudoku’ unitary, whose entries are partial symmetries; and further considering separately positive and negative part of each partial symmetry express the relations between them by placing them in a ‘double sudoku’, i.e. a magic unitary of dimension $4p$ by $4p$, which has the following block-matrix structure:

$$(7.3) \quad \begin{bmatrix} P & Q & R & S \\ Q & P & S & R \\ R & S & P & Q \\ S & R & Q & P \end{bmatrix},$$

with P, Q, R, S being p by p matrices whose entries are projections. Note that this fits with the fact that $H_s^+(p, 0)$ is a quantum subgroup of H_{2p}^+ . We can now apply Theorem 2.3 (2) of [BaV] to deduce that the algebra $A_{hs}(p, 0)$ is isomorphic to $C(H_p^{4+})$; one can check that this isomorphism is also a Hopf $*$ -algebra morphism. \square

Note that the fundamental corepresentation of $C(H_p^{4+})$ in (7.2) is in a sense intermediate between that of [BBCC] (which has dimension p and is given by the matrix with entries of the type $p_A - q_A + i(p_B - q_B)$) and that given in (7.3) considered in [BaV] (which has dimension $4p$). Once again we describe the quantum group studied in terms of the intertwiners between the tensor powers of the fundamental corepresentation.

Proposition 7.6. *The algebra $A_{hs}(p, q)$ is the universal C^* -algebra generated by the entries of a unitary $2p + q$ by $2p + q$ matrix U such that the vector ξ defined in Proposition 2.2 is a fixed vector for $U^{\otimes 2}$ and the map $e_{i\alpha} \rightarrow e_{i\alpha}^- \otimes e_{i\alpha} \otimes e_{i\alpha}^-$, $e_M \rightarrow e_M \otimes e_M \otimes e_M$ defines a morphism in $\text{Hom}(U; U^{\otimes 3})$.*

Proof. Identical to that of Proposition 3.7. \square

It remains to describe the category describing the representations of $H_s^+(p, q)$. Once again we will first separately consider the case $p = 0$ (when everything reduces to the usual computations with H_q^+ and the corresponding category is NC_{even}) and

the case $q = 0$. For the second case recall Definition 4.3 and denote by $NC_{\bullet, \text{even}}$ the subcategory given by the bulleted partitions of even size.

Theorem 7.7. *Let U denote the fundamental representation of $H_s^+(p, 0)$ introduced in Definition 7.2. We have*

$$\text{Hom}(U^{\otimes k}; U^{\otimes l}) = \text{span}(\tilde{T}_\pi \mid \pi \in NC_{\bullet, \text{even}}(k, l)),$$

where the implementing maps are defined as in (4.2).

Proof. The proof is very similar to that of Theorem 4.4. It is standard to check that $NC_{\bullet, \text{even}}$ is a tensor category satisfying the properties needed to apply the Tannaka-Krein duality of [Wor₂] and that the map $\pi \rightarrow \tilde{T}_\pi$ transforms the natural operations (concatenation, tensoring, adjoint) to the corresponding operations on the level of linear maps. Theorem 7.6 implies that the category of representations of $S^+(p, 0)$ is generated by the morphisms \tilde{T}_{π_1} and \tilde{T}_{π_2} associated with the partitions

$$\pi_1 = \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \circ \quad \circ \end{array}, \quad \pi_2 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \circ \quad \circ \end{array}.$$

The partitions π_1 and π_2 generate the whole $NC_{\bullet, \text{even}}$. Indeed, the analogous ‘non-bulleted’ partitions generate NC_{even} , and it suffices to notice that π_1 and π_2 can be composed to obtain the ‘exchange’ partition $\pi_3 = \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array}$. The generation statement is now clear.

The argument above shows that the category of the representations of $H_s^+(p, 0)$ is at least as big as $NC_{\bullet, \text{even}}$; hence there is an inclusion $\mathbb{G} \supset S^+(p, 0)$, where \mathbb{G} is the compact quantum group arising as the Tannaka-Krein dual of $NC_{\bullet, \text{even}}$ via the described implementation. To conclude the proof it suffices to show that for each $k \in \mathbb{N}$ the number $d_k := \text{card}(\text{Hom}(1; U^{\otimes 2k}))$ is equal to $\text{card}(NC_{\bullet, \text{even}}(0, 2k))$. The numbers d_k are expressed by the formula

$$d_k = h_{H_s^+(p, 0)}(\text{Tr}(U^{\otimes 2k})),$$

where $h_{H_s^+(p, 0)}$ denotes the Haar state. Using the argument similar to that of Corollary 7.4 of [BCS] and the identification in the proof of Theorem 7.5 one can show that the variables $\mathbf{p} = p_A + p_B + p_C + p_D + \dots$ and $\mathbf{q} = q_A + q_B + q_C + q_D + \dots$ (see the matrix (7.2)) are free Poisson variables of parameter $\frac{1}{4}$ – note that this provides a concrete realisation of the variables found in [BBCC]. Thus we are left with the computation of the moments of 2 copies of a difference of such two free Poisson variables. This can be deduced from the results in [BBCC]: first note that according to Theorem 7.1 of [BBCC] the $2k$ -th moment of $\mathbf{p} - \mathbf{q}$ is the k -th moment of the free Bessel law with parameters $(2, \frac{1}{2})$, and then combine the observation in the proof of Theorem 5.2 of [BBCC] with Theorem 4.3 of the same paper that the latter is equal to $\sum_{\pi \in NC_{\text{even}}(0, 2k)} (\frac{1}{2})^{\nu(\pi)}$, so that

$$(7.4) \quad d_k = 2^k \sum_{\pi \in NC_{\text{even}}(0, 2k)} \left(\frac{1}{2}\right)^{\nu(\pi)} = \sum_{\pi \in NC_{\text{even}}(0, 2k)} 2^{k-\nu(\pi)}$$

(where $\nu(\pi)$ denoted the number of blocks in the partition π). As in the proof of Theorem 4.4 we observe that the number in (7.4) indeed equals $\text{card}(NC_{\bullet, \text{even}}(0, 2k))$ due to the multiplication factor expressing the choice of colourings of bullets in each block of π . \square

We are now ready to describe the category of partitions corresponding to the representation theory of $H_s^+(p, q)$.

Theorem 7.8. *Let U denote the fundamental representation of $H_s^+(p, q)$ introduced in Definition 7.2. Then for all $k, l \in \mathbb{N}_0$*

$$\text{Hom}(U^{\otimes k}; U^{\otimes l}) = \text{span}(\hat{T}_\pi | \pi \in (NC_{\bullet, \text{even}} \star NC_{\text{even}})(k, l)).$$

Proof. We only provide the argument for the equality of the number of partitions in $(NC_{\bullet, \text{even}} \star NC_{\text{even}})(0, k)$ and the k -th moment of the character of the defining representation of $H_s^+(p, q)$. The latter can be expressed via the moment-cumulant formula (4.4), with the free cumulants $(\kappa_k)_{k=1}^\infty$ given by the sum of the free cumulants of the corresponding laws of the characters for $H_s^+(p, 0)$ and $H_s^+(0, q)$. As the second law is the free Bessel distribution, another use of the moment-cumulant formula (4.4) and Theorem 4.3 of [BBCC] implies that the even free cumulants for it are equal to 1, and the odd ones vanish. As the free Bessel laws form a free convolution semigroup, Lemma 4.7 and the proof of Theorem 7.7 imply then that the odd free cumulants related to $H_s^+(0, q)$ vanish and even ones are equal to 2^{k-1} . Hence the moment-cumulant formula for the k -th moment of the law we are interested in yields:

$$d_k = \sum_{\pi \in NC_{\text{even}}(0, k)} \prod_{b: \text{block in } \pi} (2^{\text{card}(b)-1} + 1).$$

The rest of the proof follows as in Theorem 4.8. □

Acknowledgment. Part of the work on this paper was began during the visit of the second-named author to the Université Cergy-Pontoise in April 2010. The work of the first-named author was supported by the ANR grant ‘Galoisint’.

REFERENCES

- [Ban₁] T. Banica, Quantum groups and Fuss-Catalan algebras, *Comm. Math. Phys.* **226** (2002), no. 1, 221–232.
- [Ban₂] T. Banica, Quantum automorphism groups of homogeneous graphs, *J. Funct. Anal.* **224** (2005), no. 2, 243–280.
- [BBCC] T. Banica, S. Belinschi, M. Capitaine and B. Collins, Free Bessel laws, *Canad. J. Math.*, to appear, available at *arXiv:0710.5931*.
- [BaB] T. Banica and J. Bichon, Hopf images and inner faithful representations, *Glasg. Math. J.* **52** (2010), 677–703.
- [BBC₁] T. Banica, J. Bichon and B. Collins, Quantum permutation groups: a survey, *Banach Center Publ.* **78** (2007), 13–34.
- [BBC₂] T. Banica, J. Bichon and B. Collins, The hyperoctahedral quantum group, *J. Ramanujan Math. Soc.* **22** (2007), 345–384.
- [BCS] T. Banica, S. Curran and R. Speicher, Stochastic aspects of easy quantum groups, *Probab. Theory Related Fields*, to appear, available at *arXiv:0909.0188*.
- [BaG] T. Banica and D. Goswami, Quantum isometries and noncommutative spheres, *Comm. Math. Phys.* **298** (2010), 343–356.
- [BaS] T. Banica and R. Speicher, Liberation of orthogonal Lie groups, *Adv. Math.* **222** (2009), no. 4, 1461–1501.
- [BaV] T. Banica and R. Vergnioux, Fusion rules for quantum reflection groups, *J. Noncommut. Geom.* **3** (2009), 327–359.
- [BMT] E. Bédos, G.J. Murphy and L. Tuset, Co-amenability of compact quantum groups, *J. Geom. Phys.* **40** (2001), no. 2, 130–153.

- [Bho] J. Bhowmick, Quantum isometry group of the n -tori, *Proc. Amer. Math. Soc.* **137** (2009), no. 9, 3155–3161.
- [BGS] J. Bhowmick, D. Goswami and A. Skalski, Quantum isometry groups of 0-dimensional manifolds, *Trans. AMS*, to appear, available at *arXiv:0807.4288*.
- [BhS] J. Bhowmick and A. Skalski, Quantum isometry groups of noncommutative manifolds associated to group C^* -algebras, *J. Geom. Phys.* **60** (2010), no. 10, 1474–1489.
- [BDV] J. Bichon, A. De Rijdt and S. Vaes, Ergodic coactions with large multiplicity and monoidal equivalence of quantum groups, *Comm. Math. Phys.* **262** (2006), no. 3, 703–728.
- [Gos] D. Goswami, Quantum Group of Isometries in Classical and Noncommutative Geometry, *Comm. Math. Phys.* **285** (2009), no. 1, 141–160.
- [MVD] A. Maes and A. Van Daele, Notes on compact quantum groups, *Nieuw Arch. Wisk (4)* **16** (1998), no. 1-2, 73–112.
- [NiS] A. Nica and R. Speicher, “Lectures on the combinatorics of free probability”, London Mathematical Society, Lecture Note Series, Vol. 335, Cambridge University Press, 2006.
- [Rau] S. Raum, Isomorphisms and fusion rules of orthogonal free quantum groups and their complexifications, preprint available at *arXiv:1006.2979*.
- [Spe] R. Speicher, Multiplicative functions on the lattice of noncrossing partitions and free convolution, *Math. Ann.* **298** (1994), 611–628.
- [VDW] A. Van Daele and S. Wang, Universal quantum groups, *Internat. J. Math.* **7** (1996), no. 2, 255–264.
- [Voi] D.V. Voiculescu, Addition of certain noncommuting random variables, *J. Funct. Anal.* **66** (1986), 323–346.
- [Wan₁] S. Wang, Free products of compact quantum groups. *Comm. Math. Phys.* **167** (1995), no. 3, 671–692.
- [Wan₂] S. Wang, Quantum symmetry groups of finite spaces, *Comm. Math. Phys.* **195** (1998), no. 1, 195–211.
- [Wor₁] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [Wor₂] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35–76.
- [Wor₃] S.L. Woronowicz, Compact quantum groups, in A. Connes, K. Gawedzki, and J. Zinn-Justin, editors, *Symétries Quantiques, Les Houches, Session LXIV, 1995*, 845–884.

DEPARTMENT OF MATHEMATICS, CERGY-PONTOISE UNIVERSITY, 95000 CERGY-PONTOISE, FRANCE
Email address: `teodor.banica@u-cergy.fr`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER, LA1 4YF, UNITED KINGDOM
 INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES, UL.ŚNIADECKICH 8, 00-956 WARSZAWA, POLAND
Email address: `a.skalski@lancaster.ac.uk`